

List of species deposited in the Helminthological Collection of the Institute of Parasitology, ASCR, in České Budějovice

(Abbreviations: A – Acanthocephala; C – Cestoda; Cr – Crustacea (not helminths); D – Trematoda (Digenea); L – Linguatulida (not helminths); M – Monogenea; N – Nematoda)

<i>abei</i> Asakawa, Kamiya et Ohbayashi, 1988 (<i>Soboliphyme</i>)	N – 447
<i>abnormalis</i> May, 1920 (<i>Nematodirus</i>)	N – 538
<i>abscisus</i> (Riggenbach, 1896) (<i>Choanoscolex</i>)	C – 257
<i>absconditum</i> (Looss, 1901) (<i>Acanthostomum</i>)	D – 28
<i>acanthorhodei</i> Ergens et Yukhimenko, 1975 (<i>Gyrodactylus</i>)	M – 142
<i>acanthus</i> Martin, 1975 (<i>Plethorchis</i>)	D – 501
<i>acaudatus</i> Ferraz et Thatcher, 1990 (<i>Camallanus</i>)	N – 527
<i>acanthorhyncha</i> (Wedl, 1855) Kowalewski, 1904 (<i>Tatria</i>)	C – 186
<i>acer</i> (Mueller, 1936) (<i>Pterocleidus</i>)	M – 402
<i>acipenserinum</i> Cholodkowski, 1918 (<i>Eubothrium</i>)	C – 423
<i>acipenseris</i> Wagner, 1867 (<i>Cystoopsis</i>)	N – 575
<i>acipenseri</i> Choudhury et Dick, 1992 (<i>Spinitectus</i>)	N – 659
<i>actitis</i> McIntosh, 1932 (<i>Leucochloridium</i>)	D – 141
<i>acuminata</i> (Molin, 1860) (<i>Rhabdochona</i>)	N – 111
<i>acus</i> (Bloch, 1779) (<i>Raphidascaris</i>)	N – 68
<i>acutum</i> (Linton, 1910) (<i>Schikhobalotrema</i>)	D – 358
<i>adamsoni</i> Moravec et Sey, 19 (<i>Maxvachonia</i>)	N – 449
<i>adcotylophorus</i> Manter, 1947 (<i>Opistholebes</i>)	D – 11
<i>aduncum</i> (Rudolphi, 1802) (<i>Hysterothylacium</i>)	N – 147
<i>aduncum aduncum</i> (Rudolphi, 1802) (<i>Hysterothylacium</i>)	N – 735
<i>aegyptiacus</i> Ryšavý et Moravec, 1975 (<i>Bothriocephalus</i>)	C – 14
<i>aegypticus</i> Ergens, 1981 (<i>Cichlidogyrus</i>)	M – 117
<i>aegyptiacus</i> Moravec, Baruš et Ryšavý, 1987 (<i>Skrjabinodon</i>)	N – 182
<i>aeleithrium</i> Price et Berry, 1966 (<i>Trianchoatus</i>)	M – 330
<i>aequalis</i> (Looss, 1902) (<i>Heterophyes</i>)	D – 293
<i>affine</i> Leidy, 1856 (<i>Spironoura</i>)	N – 126
<i>affinis</i> Scott, 1910 (<i>Calicotyle</i>)	M – 416
<i>affixus</i> (Looss, 1918) (<i>Pygorchis</i>)	D – 586
<i>afghana</i> Baruš, Kullmann et Tenora, 1972 (<i>Spironoura</i>)	N – 351
<i>afganensis</i> Ergens, 1979 (<i>Gyrodactylus</i>)	M – 245
<i>afghanus</i> Moravec et Amin, 1978 (<i>Neoechinorhynchus</i>)	A – 13
<i>africana</i> Moravec, Boomker et Taraschewski, 1999 (<i>Paraquimperia</i>)	N – 744
<i>africanus</i> Moravec et Van As, 2000 (<i>Philometroides</i>)	N – 764
<i>africanum</i> Moravec et Van As, 2004 (<i>Labeonema</i>)	N – 820
<i>agilis</i> (Rudolphi, 1819) (<i>Neoechinorhynchus</i>)	A – 10
<i>agonostomi</i> (<i>Creptotrema</i>) Salgado-Maldonado, Cabañas-Carranza et Caspeta-Mandujano, 1998	D – 412
<i>agonostomi</i> Moravec, Salgado-Maldonado et Caspeta-Mandujano, 1999 (<i>Paracapillaroides</i>)	N – 725
<i>agonostomi</i> Moravec et Baruš, 1971 (<i>Spinitectus</i>)	N – 106
<i>agostinhoi</i> (Pavanelli and Machado dos Santos, 1992) (<i>Chambriella</i>)	C – 402
<i>aguiari</i> Travassos, 1937 (<i>Monodontus</i>)	N – 301

<i>aguirrei</i> Travassos, 1949 (<i>Cosmoxynemoides</i>)	N – 586
<i>aguirrepequeno</i> (Jiménez, 1973) (<i>Paracreptotrematina</i>)	D – 415
<i>aharonii</i> Witenberg, 1929 (<i>Cercarioides</i>)	D – 190
<i>acheilognathi</i> Yamaguti, 1934 (<i>Bothriocephalus</i>)	C – 15
<i>achmerowi</i> Gussev, 1955 (<i>Dactylogyrus</i>)	M – 79
<i>aksuensis</i> Ergens et Karabekova, 1980 (<i>Gyrodactylus</i>)	M – 251
<i>alata</i> (Goeze, 1782) (<i>Alaria</i>)	D – 80
<i>alata</i> Rudolphi, 1819 (<i>Physaloptera</i>)	N – 232
<i>alatae</i> De et Maity, 1994 (<i>Dichelyne</i>)	N – 638
<i>alatus</i> Linstow, 1878 (<i>Dactylogyrus</i>)	M – 15
<i>alatus</i> (Ortlepp, 1932) (<i>Molinostrongylus</i>)	N – 411
<i>alatus</i> Wedl, 1861 (<i>Thelandros</i>)	N – 402
<i>alaudae</i> Rudolphi, 1819 (<i>Capillaria</i>)	N – 314
<i>albidus</i> (Braun, 1893) (<i>Metorchis</i>)	D – 410
<i>alburni</i> (Khotenovsky, 1982) (<i>Paradiplozoon</i>)	M – 301
<i>alcaraziensis</i> Lafuente et Roca, 1995 (<i>Skrjabinodon</i>)	N – 661
<i>alcoveri</i> Mas-Coma et Esteban, 1984 (<i>Tenoranema</i>)	N – 158
<i>alestesnursi</i> Paperna, 1979 (<i>Annulotrema</i>)	M – 325
<i>algericum</i> Prokopič, Hulínská et Záhř, 1983 (<i>Choniagnium</i>)	N – 98
<i>alii</i> (Rasheed, 1960) (<i>Serratospiculoides</i>)	N – 439
<i>alfortense</i> Railliet et Dollfus, 1964 (<i>Brachylentham</i>)	D – 164
<i>allaeri</i> Campana-Rouget, 1961 (<i>Spinitectus</i>)	N – 16
<i>allocytti</i> (Tkachuk, 1979) (<i>Trifoliovarium</i>)	D – 557
<i>alloyragraphorus</i> (MacCallum et MacCallum, 1913) (<i>Pyragraphorus</i>)	M – 361
<i>altavelae</i> Neifar, Euzet et Hassine, 2002 (<i>Anthobothrium</i>)	C – 369
<i>alvarengai</i> Freitas, 1957 (<i>Parapharyngodon</i>)	N – 649
<i>alveatum</i> (Creplin, 1846) (<i>Paramonostomum</i>)	D – 581
<i>amaniensis</i> (Sandground, 1928) (<i>Pseudabbreviata</i>)	N – 457
<i>amazonensis</i> Ferraz et Thatcher, 1992 (<i>Paracamallanus</i>)	N – 594
<i>amazonica</i> de Chambrier et Vaucher, 1997 (<i>Monticellia</i>)	C – 405
<i>amazonicum</i> (Travassos, 1960) (<i>Alinema</i>)	N – 831
<i>ambiguus</i> (Van Cleave, 1921) (<i>Paratenuisentis</i>)	A – 28
<i>ambiguus</i> (Rudolphi, 1819) (<i>Passalurus</i>)	N – 414
<i>ambiguus</i> (Dujardin, 1845) (<i>Proteocephalus</i>)	C – 235
<i>ambloplitis</i> (Leidy, 1887) (<i>Proteocephalus</i>)	C – 349, C – 381
<i>americana</i> Cram, 1927 (<i>Tetrameres</i>)	N – 472
<i>americana</i> Maplestone, 1930 (<i>Cruzia</i>)	N – 261
<i>americana</i> Moravec et Huffman, 2000 (<i>Dendronucleata</i>)	A – 67
<i>americana</i> Walton, 1927 (<i>Diplotriaena</i>)	N – 493
<i>americana</i> Webster, 1951 (<i>Progynotaenia</i>)	C – 174
<i>americanum</i> (Perez-Vigueras, 1956) (<i>Acanthostomum</i>)	D – 442
<i>americanum</i> Denton, 1945 (<i>Brachylecithum</i>)	D – 221
<i>americanum</i> Moravec, Borucinska et Frasca, 2002 (<i>Pancreatonema</i>)	N – 780
<i>amphibiophila</i> Moravec et Kaiser, 199 (<i>Trichospirura</i>)	N – 602
<i>amphibothrium</i> Wagener, 1857 (<i>Dactylogyrus</i>)	M – 22
<i>amplus</i> Hopper, 1967 (<i>Metoncholaimus</i>)	N – 464
<i>ampullacea</i> Miller et Harkema, 1962 (<i>Ascocotyle</i>)	D – 400
<i>amurensis</i> Achmerow, 1952 (<i>Dactylogyrus</i>)	M – 71
<i>amurensis</i> Kuperman, 1973 (<i>Triaenophorus</i>)	C – 431
<i>amurensis</i> larv. Kuperman, 1974 (<i>Triaenophorus</i>)	C – 432

<i>anabantis</i> Pearse, 1933 (<i>Camallanus</i>)	N – 393
<i>anatinum</i> Skrjabin, 1916 (<i>Epomidiostomum</i>)	N – 435
<i>anatinus</i> Markow, 1903 (<i>Prosthogonimus</i>)	D – 137
<i>anatis</i> (Schrank, 1790) (<i>Capillaria</i>)	N – 319
<i>anatis</i> Garkavi, 1965 (<i>Paramonostomum</i>)	D – 529
<i>anceps</i> (Molin, 1859) (<i>Moliniella</i>)	D – 176
<i>ancistri</i> Moravec, Gelnar et Řehulka, 1987 (<i>Capillostrongyloides</i>)	N – 175
<i>andersoni</i> Moravec, 1990 (<i>Parathelandros</i>)	N – 446
<i>androphora</i> Nitzsch, 1821 (<i>Hedruris</i>)	N – 141
<i>angorae</i> Ergens et Ibragimov, 1979 (<i>Gyrodactylus</i>)	M – 247
<i>anguillae</i> (Müller, 1980) (<i>Acanthocephalus</i>)	A – 2
<i>anguillae</i> (Leeuwenhoek, 1722) (<i>Bothriocephalus</i>)	C – 378
<i>anguillae</i> Ergens, 1960 (<i>Gyrodactylus</i>)	M – 176
<i>anguillae</i> (Yin et Sproston, 1948) (<i>Pseudodactylogyrus</i>)	M – 339
<i>anguillae</i> Moravec et Køie, 1987 (<i>Daniconema</i>)	N – 240
<i>anguillae</i> (larv.) (<i>Daniconema</i>)	N – 630
<i>anguillae</i> Moravec et Taraschewski, 2004 (<i>Procamallanus</i>)	N – 797
<i>anguillae</i> Spaul, 1927 (<i>Rhabdochona</i>)	N – 691
<i>anguillae</i> Macfarlane, 1951 (<i>Stegodexamene</i>)	D – 607
<i>angulatum</i> Linstow, 1907 (<i>Phyllodistomum</i>)	D – 122
<i>angusticolle</i> (Molin, 1860) (<i>Porrocaecum</i>)	N – 509
<i>anchoratus</i> (Dujardin, 1845) (<i>Dactylogyrus</i>)	M – 32
<i>anchoratus</i> Lim, 1987 (<i>Cornudiscoides</i>)	M – 307
<i>anisopharynx</i> Popazoglo et Boeger, 2000 (<i>Gyrodactylus</i>)	M – 358
<i>annulatus</i> Baylis et Dubney, 1922 (<i>Spiroxys</i>)	N – 522
<i>annulospinosa</i> Pichelin et Cribb, 2001 (<i>Transvena</i>)	A – 72
<i>annulosa</i> (Dujardin, 1845) (<i>Aonchotheca</i>)	N – 317
<i>anolisi</i> Baruš et Coy Otero, 1986 (<i>Oswaldocruzia</i>)	N – 376
<i>anoura</i> Dujardin, 1845 (<i>Polydelphis</i>)	N – 137
<i>anseris</i> (Madsen, 1945) (<i>Baruscapillaria</i>)	N – 327
<i>anseris</i> (Zeder, 1800) (<i>Amidostomum</i>)	N – 490
<i>anseris</i> Dubois 1961 (<i>Australapatemon</i>)	D – 536
<i>anthicum</i> Bullard et Overstreet, 2006 (<i>Psettarium</i>)	D – 537
<i>anthuris</i> (Rudolphi, 1819) (<i>Acuaria</i>)	N – 222
<i>antillarum</i> Baruš et Coy Otero, 1974 (<i>Spaulingodon</i>)	N – 346
<i>antipini</i> (Mosgovojev, 1950) (<i>Dujardinascaris</i>)	N – 239
<i>annulata</i> Moravec et Thatcher, 2001 (<i>Spinoxyuris</i>)	N – 777
<i>anudarini</i> Ergens et Dulmaa, 1968 (<i>Gyrodactylus</i>)	M – 196
<i>aphyae</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 164
<i>apophalliformis</i> Szidat, 1956 (<i>Acanthostomoides</i>)	D – 426
<i>appendiculatus</i> (Rudolphi, 1802) (<i>Hemiurus</i>)	D – 495
<i>appendiculatus</i> (Ciurea, 1916) (<i>Mesostephanus</i>)	D – 576
<i>aquaticus</i> Linnaeus, 1766 (<i>Gordius</i>)	N – 120
<i>araujo</i> Fernandes, Campos et Artigas, 1983 (<i>Travnema</i>)	N – 585
<i>araxiana</i> (Massino, 1924) (<i>Spironoura</i>)	N – 134
<i>arcuata</i> (Kowalewski, 1904) Spasskii et Spasskaja, 1954 (<i>Microsomacanthus</i>)	C – 157
<i>arcuatus</i> (Dujardin, 1845) (<i>Brachylaima</i>)	D – 524
<i>arcuatus</i> Bychowsky, 1933 (<i>Gyrodactylus</i>)	M – 191
<i>arcuatus</i> (Brandes of Stossich, 1902) (<i>Hyptiasmus</i>)	D – 129
<i>ardae</i> (Fuhrmann, 1906) (<i>Drepanidotaenia</i>)	C – 131

<i>ardeae</i> (Rausch, 1955) (<i>Dendrouterina</i>)	C – 346
<i>arenula</i> (Creplin, 1825) (<i>Lecithodolffusia</i>)	D – 572
<i>argentinensis</i> Braicovich, Moravec et Timi, 2006 (<i>Moravecica</i>)	N – 861
<i>arii</i> Moravec, Wolter et Körting, 1999 (<i>Pseudogorgorhynchus</i>)	A – 63
<i>arlenae</i> Bray et Gibson, 1995 (<i>Lepidapedon</i>)	D - 544
<i>armatum</i> (MacCallum, 1895) (<i>Homalometron</i>)	D – 200
<i>armatus</i> (Tanabe, 1922) (<i>Centrocestus</i>) mtc.	D – 551
<i>armeniaca</i> (Cholodkovski, 1915) (<i>Khawia</i>)	C – 48
<i>armstrongi</i> Gibson et Bray, 1982 (<i>Tellervotrema</i>)	D – 548
<i>artemisiana</i> Schmerling, 1925 (<i>Diplotriaena</i>)	N – 511
<i>arthracanthus</i> Paperna, 1960 (<i>Cichlidogyrus</i>)	M – 118
<i>arvicanthidis</i> (Kofed, 1915) (<i>Intermicapsifer</i>)	C – 370
<i>arvicolae</i> Schule et Skvotsov, 1931 (<i>Plagiorchis</i>)	D – 506
<i>ascaroides</i> (Goeze, 1782) (<i>Goezia</i>)	N – 27
<i>asiaticus</i> Ergens, 1978 (<i>Gyrodactylus</i>)	M – 237
<i>asoti</i> (Yamaguti, 1937) (<i>Ancylo-discoides</i>)	M – 105
<i>asperum</i> (Wright, 1879) (<i>Pegosomum</i>)	D – 583
<i>assula</i> (Dujardin, 1845) (<i>Telorchis</i>)	D – 84
<i>astorquii</i> (Watson, 1976) (<i>Acanthostomum</i>) mtc.	D – 334
<i>astyanace</i> Scholz, Aguirre-Macedo et Choudhury, 2004 (<i>Astyanace</i>)	D – 482
<i>astyanactis</i> (Watson, 1976) (<i>Genarchella</i>)	D – 312
<i>astvanactis</i> (Aatson, 1976) (<i>Genarchella</i> ?) mtc.	D – 336
<i>asymmetrica</i> Janicki, 1904 (<i>Hymenolepis</i>)	C – 274
<i>asymmetrica</i> (Janicki, 1904) (<i>Vampirolepis</i>)	C – 394
<i>atomon</i> (Rudolphi, 1802) Odhner, 1905 (<i>Podocotyle</i>)	D – 108
<i>atra</i> Nezlubinski, 1926 (<i>Tanaisia</i>)	D – 589
<i>atractostei</i> Moravec, Salgado-Maldonado, 2002 (<i>Cystoopsis</i>)	N – 792
<i>attenuata</i> (Molin, 1858) (<i>Polydelphis</i>)	N – 241
<i>attennata</i> Goldberger, 1911 (<i>Telorchis</i>)	D – 353
<i>attenuatoverrucosa</i> (Molin, 1858) (<i>Diplotriaena</i>)	N – 112
<i>attenuatum</i> (Dujardin, 1845) (<i>Brachylecithum</i>)	D – 165
<i>attenuatum</i> Mueller et Van Cleave, 1932 (<i>Lissorchis</i>)	D – 27
<i>attenuatum</i> (Dujardin, 1845) (<i>Lutztrema</i>)	D – 434
<i>attenuatum</i> (Linstow, 1906) (<i>Neodiplostomum</i>)	D – 530
<i>attenuatus</i> (Rudolphi, 1909) (<i>Notocotylus</i>)	D – 147
<i>auctum</i> (Rudolphi, 1802) (<i>Hysterothylacium</i>)	N – 737
<i>auriculatum</i> (Wedl, 1857) (<i>Crepidostomum</i>)	D – 44
<i>auriculatus</i> (Nordmann, 1832) (<i>Dactylogyrus</i>)	M – 33
<i>auriculatum</i> Yamaguti, 1952 (<i>Platybothrium</i>)	C – 276
<i>auritus</i> (Rudolphi, 1819) (<i>Glossocercus</i>) larv.	C – 281
<i>auritus</i> (Rudolphi, 1819) (<i>Glossocercus</i>)	C – 344
<i>australi</i> Vizzi et Brugni, 2003 (<i>Acolpenteron</i>)	M - 379
<i>australiensis</i> Johnston et Mawson, 1940 (<i>Anguillicola</i>)	N – 637
<i>australiensis</i> Moravec et Hobbs, 2004 (<i>Micropleura</i>)	N - 804
<i>australiensis</i> Moravec et Sey, 1990 (<i>Rhabdias</i>)	N – 450
<i>australiensis</i> Ribu et Lester, 2004 (<i>Moravecica</i>)	N - 805
<i>auziensis</i> (Seurat, 1917) (<i>Spauligodon</i>)	N – 183
<i>avitellina</i> Rego et Pavanelli, 1987 (<i>Travasssiella</i>)	C - 41
<i>avium</i> Cram, 1929 (<i>Strongyloides</i>)	N – 296
<i>awakurai</i> Ogawa et Egusa, 1978 (<i>Tetraonchus</i>)	M – 277

<i>ayalai</i> Caballero, 1971 (<i>Ascarophis</i>)	N – 695
<i>babei</i> Ha Ky, 1971 (<i>Spironoura</i>)	N – 160
<i>bacillatus</i> (Eberth, 1863) (<i>Eucoleus</i>)	N – 530
<i>baculus</i> (Diesing, 1850) (<i>Echinoparyphium</i>)	D – 144
<i>bagarii</i> Karve et Naik, 1951 (<i>Procamallanus</i>)	N – 200
<i>bagarii</i> Moravec et Sey, 1989 (<i>Dollfustrema</i>)	D – 216
<i>bagarii</i> Moravec et Sey, 1989 (<i>Cathayacanthus</i>)	A – 35
<i>bagarii</i> Moravec et Sey, 1989 (<i>Metadena</i>)	D – 223
<i>bagri</i> Lim, 1987 (<i>Cornudisoides</i>)	M – 308
<i>bagri</i> Holcman-Spector et Mane Garzon, 1988 (<i>Proteocephalus</i>)	C – 343
<i>bahamensis</i> Moravec, Coy Otero et Baruš, 1990 (<i>Abbreviata</i>)	N – 465
<i>bainae</i> (Justine et Radujković, 1988) (<i>Pseudocapillaria</i>)	N – 598
<i>bakeri</i> Moravec et Sey, 1986 (<i>Oswaldocruzia</i>)	N – 169
<i>balatonense</i> Moravec, Molnár et Székely, 1997 (<i>Lucionema</i>)	N – 689
<i>baltica</i> Szidat, 1942 (<i>Khawia</i>)	C – 42
<i>bam</i> (Tripathi, 1959) (<i>Mastacembelocleidus</i>)	M – 403
<i>banningi</i> Moravec, 1986 (<i>Huffmanella</i>)	N – 156
<i>baracoa</i> Spiridonov, 1988 (<i>Heth</i>)	N – 265
<i>barracoa</i> Baruš et Otero, 1966 (<i>Abbreviata</i>)	N – 344
<i>barbatuli</i> Ergens, 1970 (<i>Paragyrodactylus</i>)	M – 272
<i>barbatuli</i> Achmerov, 1952 (<i>Gyrodactylus</i>)	M – 184
<i>barbi</i> Ergens, 1976 (<i>Gyrodactylus</i>)	M – 223
<i>barbi</i> Baylis, 1923 (<i>Cucullanus</i>)	N – 20
<i>barbi</i> Moravec, Šimková, Hanzelová, Špakulová et Cakić, 2005 (<i>Philometroides</i>)	N – 850
<i>barusi</i> Durette- Desset, 19 (<i>Oswaldocruzia</i>)	N – 646
<i>bassii</i> (Walton, 1940) (<i>Parapharyngodon</i>)	N – 213
<i>batrachoides</i> Tubangui, 1931 (<i>Orientocreadium</i>)	D – 23
<i>baueri</i> Gussev, 1955 (<i>Dactylogyrus</i>)	M – 69
<i>baylisi</i> (Chatterji, 1935) (<i>Parapharyngodon</i>)	N – 364
<i>baylisi</i> Chabaud, 1956 (<i>Abbreviata</i>)	N – 341
<i>baylisi</i> (Travassos, 1933) (<i>Brevimulticaecum</i>)	N – 443
<i>baylisi</i> (Chabaud, 1956) (<i>Physaloptera</i>)	N – 233
<i>beema</i> (Clerc, 1906) (<i>Angularella</i>)	C – 53
<i>bequaerti</i> Viguera, 1943 (<i>Cittotaenia</i>)	C – 388
<i>beleocephalus</i> (Linstow, 1873) (<i>Echinochasmus</i>)	D – 566
<i>belidus</i> Lim et Furtado, 1986 (<i>Notopterodiscoides</i>)	M – 317
<i>benedeni</i> (Moniez, 1879) (<i>Moniezia</i>)	C – 106
<i>bengalensis</i> Nandi, De et Majumdar, 1988 (<i>Aviculariella</i>)	N – 152
<i>bernardi</i> Romashov, 1983 (<i>Eucoleus</i>)	N – 385
<i>beradae</i> Pariselle et Euzet, 2003 (<i>Cichlidogyrus</i>)	N – 385
<i>bicornis</i> Boyce, 1971 (<i>Ezonema</i>)	N – 284
<i>bicornis</i> Malewitszkaja, 1941 (<i>Dactylogyrus</i>)	M – 42
<i>bicornis</i> Moravec et Thatcher, 1996 (<i>Myleusnema</i>)	N – 657
<i>bidentatum</i> (Linstow, 1899) (<i>Hysterothylacium</i>)	N – 155
<i>bifidus</i> (Canavan, 1931) (<i>Diplotriaena</i>)	N – 494
<i>bihamuli</i> Lim et Furtado, 1986 (<i>Malayanodiscoides</i>)	M – 322
<i>bilateralis</i> (Fuhrmann, 1908) (<i>Laterorchites</i>)	C – 147
<i>bilharzi</i> (Krabbe, 1869) (<i>Variolepis</i>)	C – 134
<i>bilis</i> Braun, 1980 (<i>Metorchis</i>)	D – 253
<i>bilobus</i> (Rudolphi, 1819) (<i>Patagifer</i>)	D – 582

<i>binotati</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 290
<i>bioccai</i> Orecchia et Paggi, 1987 (<i>Cucullanus</i>)	N – 574
<i>bini</i> Kikuchi, 1929 (<i>Pseudodactylogyrus</i>)	M – 380
<i>binucleatum</i> Almeyda-Artigas, 1991 (<i>Gnathostoma</i>)	N – 640
<i>bisaccata</i> (Fuhrmann, 1906) (<i>Bisaccanthes</i>)	C – 95
<i>bisonis</i> Cram, 1925 (<i>Cooperia</i>)	N – 536
<i>biremis</i> Kowalewski, 1904 (<i>Tatria</i>)	C – 187
<i>birmani</i> Konovalov, 1967 (<i>Gyrodactylus</i>)	M – 234
<i>biwakoensis</i> (<i>Raphidascaris</i>) larv.	N – 714
<i>biwakoensis</i> Fujita, 1928 (<i>Raphidascaris</i>)	N – 786
<i>blennii</i> Paggi et Orecchia, 1975 (<i>Tyrrhenia</i>)	D – 261
<i>bliccae</i> (Reichenbach-Klinke, 1961) (<i>Paradiplozoon</i>)	M – 300
<i>bliccensis</i> Gläser, 1974 (<i>Gyrodactylus</i>)	M – 230
<i>bobica</i> (Clerc, 1903) (<i>Orthoskrjabinia</i>)	C – 165
<i>bobica</i> Clerc, 1903 (<i>Anonchotaenia</i>)	C – 60
<i>bohemicum</i> Moravec, 1977 (<i>Esocinema</i>)	N – 110
<i>bolonensis</i> Ergens et Yukhimenko, 1975 (<i>Gyrodactylus</i>)	M – 222
<i>bolschewense</i> (Kotova, 1939) (<i>Echinostoma</i>)	D – 288
<i>bonneti</i> de Chambrier, Coquille et Brooks, 2006 (<i>Ophiotaenia</i>)	C – 400
<i>borealis</i> Nybelin, 1936 (<i>Dactylogyrus</i>)	M – 52
<i>borealis</i> Keeny et Campbell, 2001 (<i>Grillotia</i>)	C – 336
<i>borealis</i> (Olsson, 1893) (<i>Tetraonchus</i>)	M – 265
<i>botulovagina</i> (Gussev et Strelkow, 1960) (<i>Ancylo-discoides</i>)	M – 106
<i>bovieri</i> Blanchard, 1886 (<i>Rictularia</i>)	N – 413
<i>bozkovi</i> Moravec, 1966 (<i>Hexametra</i>)	N – 100
<i>brachycollis</i> Janiszewska, 1953 (<i>Caryophyllaeus</i>)	C – 51
<i>brachyarthera</i> Cholodkovsky, 1906 (<i>Dilepis</i>)	C – 92
<i>brachymystacis</i> Ergens, 1978 (<i>Gyrodactylus</i>)	M – 236
<i>bramae</i> (Müller, 1776) (<i>Sphaerostoma</i>)	D – 46
<i>branchialis</i> Justine, 2004 (<i>Huffmanella</i>)	N – 816
<i>branchicus</i> Malmberg, 1964 (<i>Gyrodactylus</i>)	M – 244
<i>brasiliense</i> Moravec et Thatcher, 1999 (<i>Myleusnema</i>)	N – 741
<i>brasiliensis</i> Moravec, Kohn, Fernandes, 1991 (<i>Ichthyouris</i>)	N – 548
<i>brasiliensis</i> Moravec, Kohn et Fernandes, 1995 (<i>Goezia</i>)	N – 624
<i>brasiliensis</i> Moravec et thatcher, 1997 (<i>Raphidascaroides</i>)	N – 686
<i>bravohollisae</i> Kritsky, Vidal-Martínez et Rodríguez-Canul, 1994 (<i>Sciadicleithrum</i>)	M – 382
<i>brayi</i> Kuchta et Scholz, 2004 (<i>Bathycestus</i>)	C - 303
<i>brevicaeca</i> Moravec, Kohn et Fernandes, 1995 (<i>Goezia</i>)	N – 625
<i>brevicaecum</i> Ochi in Goto et Ozaki, 1929 (<i>Mesocoelium</i>)	D – 301
<i>brevicauda</i> Boulenger, 1916 (<i>Triodontophorus</i>)	N – 544
<i>brevicaudata</i> (Rodhain et Vuylsteke, 1937) (<i>Oswaldofilaria</i>)	N – 426
<i>brevicaudatum</i> (Zeder, 1800) (<i>Oxysomatium</i>)	N – 85
<i>brevicole</i> (Creplin, 1829) (<i>Psilostomum</i>)	D – 609
<i>breviplexus</i> (Stafford, 1902) (<i>Pneumonoeces</i>)	D – 243
<i>brevis</i> Rybicka, 1957 (<i>Diorchis</i>)	C – 94
<i>brevis</i> (Clerc, 1902) (<i>Anomotaenia</i>)	C – 62
<i>brevis</i> Kirschenblat, 1938 (<i>Paranoplocephala</i>)	C – 202
<i>brevis</i> (Van Cleave, 1916) (<i>Polymorphus</i>) larv.	A – 56
<i>brevispicula</i> Moravec et Kaiser, 1994 (<i>Oswaldocruzia</i>)	N – 620

<i>brevispiculus</i> Moravec, Kohn et Fernandes, 1993 (<i>Cucullanus</i>)	N – 615
<i>bridgeri</i> Bray et Gibson, 1986 (<i>Panopula</i>)	D – 549
<i>brooksi</i> García-Prieto, Rodríguez et Pérez-Poncede León, 1996 (<i>Proteocephalus</i>)	C – 252
<i>brunensis</i> Řehulková et Gelnar, 2005 (<i>Thylacicleidus</i>)	M – 399
<i>buccalis</i> Pence et Little, 1972 (<i>Anatrichosoma</i>)	N – 258
<i>buccalis</i> (Yamaguti, 1943) (<i>Aonchotheca</i>)	N – 167
<i>buckleyi</i> (Dissanaïke, 1961) (<i>Brugia</i>)	N – 105
<i>bufonis</i> (Schrank, 1788) (<i>Rhabdias</i>)	N – 93
<i>bufonis</i> (Shiple, 1903) (<i>Pseudoacanthocephalus</i>)	A – 38
<i>bufonis</i> Vigneras 1942 (<i>Batrachotaenia</i>)	C – 224
<i>bursatum</i> Spiridonov, 1988 (<i>Ransomnemnema</i>)	N – 267
<i>bulbosus</i> (Linstow, 1899) (<i>Parapharyngodon</i>)	N – 184
<i>bullatatarudis</i> Turnbull, 1956 (<i>Gyrodactylus</i>)	M – 364
<i>bului</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 291
<i>bungara</i> Srivastava et Ghosh, 1968 (<i>Encyclometra</i>)	D – 281
<i>bungara</i> Srivastava et Ghosh, 1968 (<i>Encyclometra</i>) mtc.	D – 299
<i>buongerminii</i> Lent, Freitas et Cavalcanti, 1945 (<i>Prosthodendrium</i>)	D – 161
<i>bursata</i> Moravec et Prouza, 1995 (<i>Ichthyouris</i>)	N – 653
<i>bursicola</i> Odhner, 1900 (<i>Gymnophallus</i>)	D – 592
<i>bursicola</i> (Creplin, 1837) (<i>Uroproctepisthmium</i>)	D – 590
<i>buschkieli</i> Bychowsky, 1957 (<i>Heteronchocleidus</i>)	M – 259
<i>buteonis</i> (Schrank, 1788) (<i>Centrorhynchus</i>)	A – 68
<i>bychowskyi</i> Caballero et Caballero, 1975 (<i>Perezitrema</i>)	D – 465
<i>caballeroi</i> (Peláez et Cruz, 1953) (<i>Acanthostomum</i>)	D – 443
<i>caballeroi</i> Ryšavý et Macko, 1973 (<i>Paradilepis</i>) larv.	C – 313
<i>caballeroi</i> Ryšavý et Macko, 1973 (<i>Paradilepis</i>)	C – 345
<i>cableri</i> Martin et Zam, 1967 (<i>Vasorhabdochona</i>)	N – 724
<i>caballeroi</i> Pritchard, 1960 (<i>Preptetos</i>)	D – 347
<i>caballeroi</i> Baruš et Coy Otero, 1966 (<i>Terranova</i>)	N – 483
<i>cahirinus</i> (Looss, 1896) (<i>Haplorchoides</i>)	D – 34
<i>calculus</i> Looss, 1907 (<i>Pachytrema</i>)	D – 134
<i>calicatum</i> (Looss, 1900) (<i>Trichonema</i>)	N – 332
<i>californiense</i> Moravec, Spangenberg, Slager, Lewand, Sim et Frasca, 2001 (<i>Syngnathinema</i>)	N – 770
<i>californiensis</i> (Cort, 1919) (<i>Glyphelmis</i>)	D – 67
<i>californiensis</i> Martin, 1950 (<i>Euhaplorchis</i>)	D – 513
<i>caligorum</i> Johnston, 1835 (<i>Udonella</i>)	M – 376
<i>cameroni</i> Rausch, 1969 (<i>Diphyllobothrium</i>)	C – 329
<i>campechensis</i> González-Solís et Moravec, 2004 (<i>Orientattractis</i>)	N – 806
<i>campylancristrota</i> (Wedl, 1855) (<i>Valipora</i>)	C – 27
<i>canadensis</i> Moravec, Conboy et Speare, 2005 (<i>Huffmanella</i>)	N – 823
<i>canadensis</i> Moravec et Arai, 1971 (<i>Rhabdochona</i>)	N – 4
<i>canadensis balsasensis</i> Caspeta, Moravec et Salgado, 1998 (<i>Rhabdochona</i>)	N – 696
<i>canadensis bifilamentosa</i> Moravec et Huffman, 1988 (<i>Rhabdochona</i>)	N – 282
<i>candzei</i> (Fraipont, 1882) (<i>Foleyella</i>)	N – 132
<i>caninum</i> Ercolani, 1859 (<i>Ancylostoma</i>)	N – 313
<i>caninum</i> (<i>Dipylidium</i>)	C – 237
<i>carolini</i> Holl, 1928 (<i>Spinitectus</i>)	N – 262

<i>cantonensis</i> Chen, 1935 (<i>Angiostrongylus</i>)	N – 260
<i>capillaroides</i> Fuhrmann, 1906 (<i>Hymenolepis</i>)	C – 135
<i>capilliformis</i> Oschmarin, 1952 (<i>Brachylecthum</i>)	D – 517
<i>capitellata</i> (Schneider, 1866) (<i>Cyrnea</i>)	N – 487
<i>capitellatum</i> (Rudolphi, 1819) (<i>Anisocoelium</i>)	D – 241
<i>capito</i> (Rudolphi, 1819) (<i>Cyclusteria</i>)	C – 278
<i>capito</i> (Rudolphi, 1819) (<i>Cyclusteria</i>) larv.	C – 279
<i>capoetai</i> Ergens et Ibragimov, 1976 (<i>Gyrodactylus</i>)	M – 225
<i>capromydis</i> Baruš et Ryšavý, 1967 (<i>Trichotravassosia</i>)	N – 484
<i>capurroi</i> Vidal-Martínez et Mendoza-Franco, 1998 (<i>Pseudorhabdosynochus</i>)	M – 350
<i>carangis</i> Olsen, 1954 (<i>Camallanus</i>)	N – 859
<i>carassii</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 147
<i>carbonis</i> (Dubinin et Dubinina, 1940) (<i>Baruscapillaria</i>)	N – 768
<i>carcharhini</i> Moravec et Little, 1988 (<i>Granulinema</i>)	N – 277
<i>caribaensis</i> (Ryšavý et Macko, 1973) (<i>Glossocercus</i>) larv.	C – 282
<i>caribaensis</i> (Ryšavý et Macko, 1973) (<i>Glossocercus</i>)	C – 418
<i>carioca</i> Freitas et Lent, 1935 (<i>Capillaria</i>)	N – 656
<i>carnosum</i> (Rudolphi, 1819) (<i>Pachycreadium</i>)	D – 264
<i>carolinensis</i> Moravec, de Buron et Roumillat, 2006 (<i>Philometra</i>)	N – 855
<i>carparum</i> (Odening, 1959) (<i>Allocreadium</i>)	D – 41
<i>carpathicus</i> Zachvatkin, 1951 (<i>Dactylogyrus</i>)	M – 34
<i>cascadilla</i> Wigdor, 1918 (<i>Rhabdochona</i>)	N – 43
<i>catenata</i> Van Cleave et Muller, 1932 (<i>Capillaria</i>)	N – 273
<i>catinatum</i> (Looss, 1900) (<i>Trichonema</i>)	N – 333
<i>catostomi</i> Cooper, 1920 (<i>Glaridacris</i>)	C – 5
<i>catulinus</i> Kostylew, 1927 (<i>Macracanthorhynchus</i>)	A – 47
<i>caudatus</i> Moravec, Scholz et Vivas Rodríguez, 1995 (<i>Neophilometroides</i>)	N – 658
<i>celatus</i> (Van Cleave, 1928) (<i>Pallisentis</i>)	A – 31
<i>cenaticum</i> (Bruce et Cannon, 1989) (<i>Hysterothylacium</i>)	N – 837
<i>cenotae</i> (Pearse, 1936) (<i>Hysterothylacium</i>)	N – 660
<i>cenotae</i> (Pearse, 1936) (<i>Hysterothylacium</i>) larv.	N – 668
<i>centripunctata</i> (Rivolta, 1974) (<i>Avitelina</i>)	C – 216
<i>centuri</i> Baruš, 1966 (<i>Tetrameres</i>)	N – 369
<i>centuri</i> Rysavy, 1966 (<i>Choanotaenia</i>)	C – 141
<i>cernuae</i> (Gmelin, 1790) (<i>Proteocephalus</i>)	C – 47
<i>cernuae</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 171
<i>cervi</i> Cameron, 1931 (<i>Elaphostrongylus</i>)	N – 381
<i>cesticillus</i> (Molin, 1858) (<i>Raillietina</i>)	C – 175
<i>cestoides</i> (Beneden, 1871) (<i>Otodistomum</i>)	D – 68
<i>cetaceum</i> Johnston et Best, 1943 (<i>Polymorphus</i>)	A – 70
<i>chabaudi</i> Baruš, 1965 (<i>Oxyspirura</i>)	N – 360
<i>chaenogobii</i> Yamaguti, 1939 (<i>Nippotaenia</i>)	C – 338
<i>chamaeleonis</i> Baylis et Daubney, 1923 (<i>Strongyluris</i>)	N – 525
<i>chamelensis</i> Pérez-Ponce de León, Brooks et Berman, 1995 (<i>Proteocephalus</i>)	C – 310
<i>chandleri</i> Lumsden, 1963 (<i>Ascocotyle</i>) mtc.	D – 352
<i>chandleri</i> Bullock, 1957 (<i>Octospiniferoides</i>)	A – 55
<i>chauhani</i> Venkatanarsaiah et Kulkarni, 1990 (<i>Chauhanellus</i>)	M – 335
<i>chavarriai</i> (Price, 1938) (<i>Ameloblastella</i>)	M – 354
<i>cheilancristrotus</i> (Wedl, 1855) (<i>Neogryporhynchus</i>)	C – 205
<i>cheligenitalis</i> Lim et Furtado, 1984 (<i>Dactylogyrus</i>)	M – 372

<i>chelydrae</i> (Harwood, 1932) Mackin, 1936 (<i>Spironoura</i>)	N – 269
<i>chetumalensis</i> González-Solís, Moravec et Vidal-Martínez, 2002 (<i>Procamallanus</i>)	N – 785
<i>chevreuxi</i> Seurat, 1918 (<i>Amidostomum</i>)	N – 521
<i>chiapasensis</i> González-Solís et Moravec 2004 (<i>Orientattractis</i>)	N – 807
<i>chimusensis</i> (Freitas et Ibáñez, 1968) (<i>Procamallanus</i>)	N – 815
“ <i>chinensis</i> “ Groschaft (?) (<i>Lyperosomum</i>)	D – 173
<i>chitwoodae</i> (Le-Van-Hoa et Pham-Ngoc-Khue, 1971) (<i>Oceanicucullanus</i>)	N – 558
<i>chodukini</i> Osmanov, 1957 (<i>Rhabdochona</i>)	N – 36
<i>chondrostomi</i> Malewitszkaja, 1941 (<i>Dactylogyrus</i>)	M – 18
<i>chondrostomi</i> Ergens, 1967 (<i>Gyrodactylus</i>)	M – 192
<i>chranilowi</i> Bychowsky, 1931 (<i>Dactylogyrus</i>)	M – 51
<i>chrysanthemoides</i> Skrjabin, 1916 (<i>Orneoscaris</i>)	N – 455
<i>cichlasomae</i> Moravec, Scholz et Mendoza-Franco, 1995 (<i>Capillaria</i>)	N – 641, N - 664
<i>cichlasomae</i> Manter, 1936 (<i>Crassicutis</i>) mtc.	D – 362
<i>cichlasomae</i> Moravec, Vidal et Salgado-Maldonado, 1992 (<i>Mexiconema</i>)	N – 573
<i>cichlidarum</i> Paperna, 1968 (<i>Gyrodactylus</i>)	M – 406
<i>cichlidorum</i> Aguirre-Macedo et Scholz (<i>Culuwiya</i>)	D – 511
<i>cingulifera</i> (Krabbe, 1869) (<i>Choanotaenia</i>)	C – 142
<i>cingulifera</i> (Krabbe, 1869) (<i>Anomotaenia</i>)	C – 65
<i>cichlasomae</i> Moravec et Baruš, 1971 (<i>Spectatus</i>)	N – 107
<i>cichlasomae</i> Manter, 1936 (<i>Crassicutis</i>)	D – 15
<i>ciliata</i> Fuhrmann, 1913 (<i>Unciunia</i>)	C – 103
<i>ciliata</i> (Fuhrmann, 1913) (<i>Anomotaenia</i>)	C – 64
<i>cinctum</i> (Rudolphi, 1803) (<i>Echinoparyphium</i>)	D – 148
<i>cinctus</i> (Linstow, 1897) (<i>Parapharyngodon</i>)	N – 343, N - 534
<i>cirratus</i> Paperna, 1964 (<i>Cichlidogyrus</i>)	M – 119
<i>citharini</i> Khalil, 1964 (<i>Cithariniella</i>)	N – 18
<i>citrus</i> (Krabbe, 1869) (<i>Anomotaenia</i>)	C – 66
<i>clariadis</i> Paperna, 1961 (<i>Quadriacanthus</i>)	M – 262
<i>clariasi</i> Dajia, Dingke et Xiaoming, 1986 (<i>Phyllodistomum</i>)	D – 211
<i>clathratum</i> (Deslongchamps, 1824) (<i>Zonorchis</i>)	D – 432
<i>clausa</i> (Rudolphi, 1819) (<i>Physaloptera</i>)	N – 140
<i>clavaiceps</i> Dogiel et Akhmerov, 1959 (<i>Philometra</i>)	N - 84
<i>clavata</i> (Nordmann, 1832) (<i>Tylodelphys</i>)	D – 87
<i>clavatus</i> Van Cleave, 1940 (<i>Gorgorhynchus</i>)	A – 51
<i>claviceps</i> (Goeze, 1782) (<i>Bothriocephalus</i>)	C – 16
<i>claviformis</i> (Brandes, 1888) (<i>Microphallus</i>)	D – 606
<i>clavula</i> (Dujardin, 1845) (<i>Acanthocephalus</i>)	A – 46
<i>clerci</i> Mathevossian et Krotov, 1949 (<i>Echinocotyle</i>)	C – 132
<i>clerci</i> Skrjabin, 1915 (<i>Echinoparyphium</i>)	D – 568
<i>cloacicola</i> (Lühe, 1909) (<i>Paralepoderma</i>)	D – 82
<i>cloaciterus</i> (Oswaria, 1956) (<i>Microtetrameres</i>)	N – 502
<i>coartatus</i> Řehulková et Gelnar, 2006 (<i>Dactylogyrus</i>)	M – 409
<i>coaxatus</i> Dietz, 1909 (<i>Echinochasmus</i>)	D – 567
<i>cobitis</i> Bychowsky, 1933 (<i>Gyrodactylus</i>)	M – 180
<i>cobitis</i> Ergens, 1963 (<i>Ancyrocephalus</i>)	M – 99
<i>cobitidis proterhorhini</i> Vojtek, 1964 (<i>Apatemon</i>)	D – 114
<i>coelica</i> Nybelin, 1941 (<i>Diplocotyle</i>)	M - 395

<i>cochleavagina</i> Gussev et Strelkow, 1960 (<i>Ancylo-discoides</i>)	M – 107
<i>cochlearii</i> Coil, 1955 (<i>Parvitaenia</i>)	C – 254
<i>colorni</i> Chisholm, Whittington et Kearns, 2001 (<i>Dendromonocotyle</i>)	M – 365
<i>collaris</i> (Batsch, 1786) Lopez-Neyra, 1942 (<i>Myxolepis</i>)	C – 160
<i>collata</i> Chatterji, 1933 (<i>Masenia</i>)	D – 215
<i>columbae</i> (Gmelin, 1970) (<i>Ascaridia</i>)	N – 226
<i>columbrimurorum</i> (Rudolphi, 1819) (<i>Encyclometra</i>)	D – 83
<i>colubrimurorum</i> (Rudolphi, 1819) (<i>Encyclometra</i>) mtc.	D – 277
<i>colymbi</i> Zeder, 1803 (<i>Ligula</i>)	C – 151
<i>commutatum</i> (Diesing, 1850) (<i>Diplostomum</i>) mtc.	D – 468
<i>commutatum</i> (Diesing, 1850) (<i>Diplostomum</i>)	D – 598
<i>compacta</i> (Clerc, 1906) (<i>Raillietina</i>)	C – 176
<i>compactum</i> (Lutz, 1928) (<i>Diplostomum</i>) mtc.	D – 321
<i>complanatum</i> (Rudolphi, 1819) (<i>Clinostomum</i>) mtc.	D – 65
<i>complanatum</i> (Rudolphi, 1814) (<i>Clinostomum</i>)	D – 357
<i>compressa</i> (Linton, 1892) (<i>Microsomacanthus</i>)	C – 153
<i>concaevum</i> (Creplin, 1825) (<i>Cryptocotyle</i>)	D – 471
<i>confertus</i> Machalska, 1974 (<i>Psilotornus</i>)	D – 612
<i>confusa</i> Hunter, 1927 (<i>Glaridacris</i>)	C – 363
<i>confusus</i> (Looss, 1894) (<i>Prosotocus</i>)	D – 91
<i>congolensis</i> Campana-Rouget, 1961 (<i>Rhabdochona</i>)	N – 35
<i>conica</i> (Fuhrmann, 1908) (<i>Orthoskrjabinina</i>)	C – 167
<i>conicum</i> Dubois, 1937 (<i>Neodiplostomum</i>)	D – 593
<i>conoidem</i> (Bloch, 1782) (<i>Hypoderaeum</i>)	D – 138
<i>conostomum</i> (Olssen, 1876) (<i>Phyllodistomum</i>)	D – 286
<i>constantinae</i> (Mueller, 1935) (<i>Leucochloridomorpha</i>)	D – 145
<i>constricta</i> (Molin, 1858) (<i>Anomotaenia</i>)	D – 63
<i>contorta</i> (Creplin, 1939) (<i>Capillaria</i>)	N – 516
<i>contortus</i> (Rudolphi, 1819) (<i>Spiroxys</i>)	N – 99
<i>contortus</i> (Rudolphi, 1803) (<i>Haemonchus</i>)	N – 452
<i>contortus</i> (Creplin, 1839) (<i>Eucoleus</i>)	N – 330
<i>contrarmatus</i> Lim et Furtado, 1984 (<i>Dactylogyrus</i>)	M – 373
<i>convolvagensis</i> Lim et Furtado, 1984 (<i>Dactylogyrus</i>)	M – 375
<i>convolvuligenitalis</i> Lim et Furtado, 1984 (<i>Dactylogyrus</i>)	M – 291
<i>cophixali</i> Moravec et Sey, 1989 (<i>Opisthiohlyphe</i>)	D – 232
<i>coraciae</i> (Gmelin, 1790) (<i>Squamofilaria</i>)	N – 442
<i>corax</i> Schell, 1953 (<i>Microtetrameres</i>)	N – 354
<i>corderoi</i> Torres, Teuber et Miranda, 1990 (<i>Camallanus</i>)	N – 605
<i>cordonspinosa</i> Baruš et Garrido, 1968 (<i>Acuaria</i>)	N – 289
<i>cordus</i> Nybelin, 1937 (<i>Dactylogyrus</i>)	M – 30
<i>cornoides</i> Gläser et Gussev, 1971 (<i>Dactylogyrus</i>)	M – 26
<i>cornutum</i> (Stossich, 1904) (<i>Hysterothylacium</i>)	N – 736
<i>coronacauda</i> Belouss, 1965 (<i>Rhabdochona</i>)	N – 41
<i>coronata</i> (Rudolphi, 1809) (<i>Squamofillaria</i>)	N – 503
<i>coronatum</i> (Looss, 1900) (<i>Trichonema</i>)	N – 334
<i>coronatus</i> Molin, 1861 (<i>Histiostrongylus</i>)	N – 217, N - 507
<i>coronula</i> (Dujardin, 1845) (<i>Dicranotaenia</i>)	C – 86
<i>cornu</i> (Zeder, 1800) (<i>Apharyngostrigea</i>)	D – 132
<i>cornu</i> Linstow, 1878 (<i>Dactylogyrus</i>)	M – 23
<i>cornutus</i> (Rudolphi, 1808) (<i>Cotylurus</i>)	D - 194

<i>corvorum</i> (Rudolphi, 1819) (<i>Pseudocapillaria</i>)	N – 325, N – 499
<i>coryphaenae</i> Bosc, 1797 (<i>Tentacularia</i>)	C – 12
<i>coryphaenae</i> Bosc, 1797 (<i>Tentacularia</i>)	C – 318
<i>costatae</i> Ergens et Gussev, 1976 (<i>Gyrodactylus</i>)	M – 220
<i>costatus alsophisi</i> Baruš et Coy Otero, 1966 (<i>Kalicephalus</i>)	N – 482
<i>costellatum</i> (Dujardin, 1845) (<i>Heligmosomum</i>)	N – 219
<i>cotti</i> Fujita, 1927 (<i>Camallanus</i>)	N – 11
<i>cotti</i> Gustafson, 1940 (<i>Rhabdochona</i>)	N – 31
<i>cotti</i> Roman, 1956 (<i>Gyrodactylus</i>)	M – 181
<i>cotylophorum</i> (Fischoeder, 1901) (<i>Cotylophoron</i>)	N – 247
<i>crami</i> Viguera, 1934 (<i>Ornithostrongylus</i>)	N – 379
<i>crassa</i> Kuwahara, Niimi et Itagaki, 1974 (<i>Anguillicola</i>)	N – 197
<i>crassicauda</i> (Bellingham, 1840) (<i>Trichosomoides</i>)	N – 708
<i>crassiusculus</i> (Rudolphi, 1809) (<i>Metorchis</i>)	D – 577
<i>crassum</i> (Bloch, 1779) (<i>Eubothrium</i>)	C – 271
<i>crassum</i> (Deslongchamps, 1824) (<i>Porrocaecum</i>)	N – 198
<i>crassus</i> Kulwiec, 1927 (<i>Dactylogyrus</i>)	M – 44
<i>crassus</i> Van Cleave, 1919 (<i>Neoechinorhynchus</i>)	A – 9
<i>crassus</i> Forel, 1868 (<i>Triaenophorus</i>)	C – 326
<i>crassus</i> Forel, 1868 (<i>Triaenophorus</i>) larv.	C – 350
<i>crassicaudata</i> (Creplin, 1829) (<i>Streptocara</i>)	N – 71
<i>crassiscolex</i> (Linstow, 1890) (<i>Choanotaenia</i>)	C – 196
<i>crateriformes</i> (Goeze, 1782) (<i>Choanotaenia</i>)	C – 143
<i>creplini</i> (Krabbe, 1869) (<i>Wardium</i>)	C – 105
<i>crenata</i> (Goeze, 1789) (<i>Variolepis</i>)	C – 191
<i>crenata</i> (Goeze, 1782) (<i>Passerilepis</i>)	C – 194
<i>crenata</i> (Goeze, 1782) (<i>Dicranotaenia</i>)	C – 87
<i>crescenticleithrium</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 293
<i>créseyi</i> Timi et Sardela, 1998 (<i>Nothobomolochus</i>)	Cr – 3
<i>cricetorum</i> Kirschenolatt, 1949 (<i>Catenotaenia</i>)	C – 287
<i>cricosoure</i> Baruš et Coy Otero, 1974 (<i>Skrjabinodon</i>)	N – 348
<i>cristatocleithrium</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 296
<i>crossophorum</i> Viguera, 1942 (<i>Mesocoelium</i>)	D – 70
<i>crotophaga</i> Yorke et Maplestone, 1926 (<i>Torquatella</i>)	N – 434
<i>cruciatus</i> (Wedl, 1857) (<i>Ancyrocephalus</i>)	M – 100
<i>cruciata</i> (Rudolphi, 1819) (<i>Raillietina</i>)	C – 177
<i>crucibulum</i> (Rudolphi, 1819) (<i>Prosochynchus</i>)	D – 259
<i>crucifer</i> Wagener, 1857 (<i>Dactylogyrus</i>)	M – 24
<i>crumenifer</i> (Creplin, 1847) (<i>Gastrothylax</i>)	D – 249
<i>cryptomeres</i> Bychowski, 1934 (<i>Dactylogyrus</i>)	M – 25
<i>ctenoides</i> (Riehm, 1881) (<i>Ctenotaenia</i>)	C – 108
<i>ctenosauri</i> Caballero, 1938 (<i>Ozolaimus</i>)	N – 647
<i>cubana</i> Baruš, 1969 (<i>Dudekemia</i>)	N – 375
<i>cubana</i> Baruš, 1972 (<i>Aplectana</i>)	N – 356
<i>cubense</i> Baruš et Sonin, 1971 (<i>Hastospiculum</i>)	N – 467
<i>cubensis</i> Baruš et Otero, 1969 (<i>Parapharyngodon</i>)	N – 345
<i>cubensis</i> Groschaft, 1965 (?) (<i>Parabascus</i>)	D – 163
<i>cubensis</i> Spiridonov, 1988 (<i>Ichthyocephalus</i>)	N – 268
<i>cubensis</i> Moravec et Coy Otero, 1987 (<i>Rhabdochona</i>)	N – 117
<i>cubensis</i> (Read et Amrein, 1953) (<i>Spaulingodon</i>)	N – 331

<i>cucullatus</i> (Wehr, 1934) (<i>Victorocara</i>)	N – 362
<i>cuniculi</i> (Blanchard, 1891) (<i>Andrya</i>)	C – 389
<i>curvicirrus</i> Achmerow, 1952 (<i>Dactylogyrus</i>)	M – 74
<i>curvilamellis f. obscura</i> Gussev et Strelkow, 1960 (<i>Ancylodiscoides</i>)	M – 108
<i>curvilamellis f. typica</i> Gussev et Strelkow, 1960 (<i>Ancylodiscoides</i>)	M – 109
<i>cuspidatus</i> Cooper, 1917 (<i>Bothriocephalus</i>)	C – 269
<i>cuspidatus</i> (Loos, 1896) (<i>Centrocestus</i>)	D – 19
<i>cuticola</i> (Nordmann, 1832) (<i>Posthodiplostomum</i>)	D – 115
<i>cyathopharyns</i> (Baylis, 1923) (<i>Paracamallanus</i>)	N – 12
<i>cycloancistrioides</i> Kritsky, Boe et Thatcher, 1985 (<i>Dawestrema</i>)	M – 421
<i>cycloancistrum</i> Price et Nowlin, 1967 (<i>Dawestrema</i>)	M – 420
<i>cygnoides</i> (Zeder, 1800) (<i>Gorgodera</i>)	D – 57
<i>cylindracea</i> (Zeder, 1800) (<i>Haplometra</i>)	D – 73
<i>cylindracea</i> (Bloch, 1782) (<i>Cladotaenia</i>)	C – 81
<i>cylindraceum</i> (Goere, 1782) (<i>Prosthorhynchus</i>)	A – 75
<i>cylindratus</i> (Van Cleave, 1913) (<i>Neoechinorhynchus</i>)	A – 12
<i>cylindrica</i> Linstow, 1883 (<i>Aprocta</i>)	N – 773
<i>cylindrica</i> (Eberth, 1863) (<i>Baruscapillaria</i>)	N – 243, N - 518
<i>cynoscicola</i> Timi et Etchegoin, 1996 (<i>Lernanthropus</i>)	Cr – 1
<i>cynoscionis</i> Moravec, de Buron et Roumillat, 2006 (<i>Philometra</i>)	N – 856
<i>cyprinacea</i> Linnaeus, 1758 (<i>Lernaea</i>)	Cr – 4
<i>cyprini</i> Diarova, 1964 (<i>Gyrodactylus</i>)	M – 188
<i>cyprini</i> Yamaguti, 1941 (<i>Cucullanus</i>)	N – 21
<i>cyprini</i> (Ishii, 1931) (<i>Philometroides</i>)	N – 572
<i>cyprini</i> Molnár et Moravec, 1997 (<i>Skrjabillanus</i>)	N – 687
<i>cyprinirutili</i> (Creplin, 1825) (<i>Philometra</i>)	N – 83
<i>cyprinodontis</i> Chandler, 1935 (<i>Glossocercus</i>)	C – 419
<i>dagestanica</i> (Altaev, 1903) (<i>Spiculopteroides</i>)	N – 177
<i>decacantha</i> Fuhrmann, 1913 (<i>Tatria</i>)	C – 188
<i>decipiens</i> Krabbe, 1878 (<i>Phocanema</i>)	N – 172
<i>decipiens</i> (Krabbe, 1878) (<i>Pseudoterranova</i>)	N – 569
<i>decorata</i> Cram, 1927 (<i>Syncuaria</i>)	N – 495
<i>decorus</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 314
<i>dehiscens</i> (Krabbe, 1879) (<i>Anomotaenia</i>)	C – 67
<i>delafondi</i> (Railliet, 1892) (<i>Aporina</i>)	C – 77
<i>delectans</i> (Braun, 1901) (<i>Zonorchis</i>)	D – 155
<i>deliciosus</i> (Olsson, 1893) (<i>Gymnophallus</i>)	D – 427
<i>delta</i> Paperna, 1979 (<i>Annulotrema</i>)	M – 326
<i>dendritica</i> (Goeze, 1782) (<i>Catenotaenia</i>)	C – 437
<i>dendriticum</i> Rudolphi, 1819 (<i>Dicrocoelium</i>)	D – 306
<i>dendriticum</i> (Nitzsch, 1824) larv. (<i>Diphyllobothrium</i>)	C – 128
<i>dentatum</i> (Rudolphi, 1803) (<i>Ornithostrongylus</i>)	N – 228
<i>dentatum</i> (Linton, 1900) (<i>Stephanostomum</i>)	D – 371
<i>dentatus</i> Moravec et Thatcher, 1997 (<i>Procamallanus</i>)	N – 678
<i>denticulata</i> (Olsson, 1876) (<i>Diclidophora</i>)	M - 401
<i>denticulatus</i> (Rudolphi, 1802) (<i>Mesorchis</i>)	D – 479
<i>denudata</i> (Dujardin, 1845) (<i>Rhabdochona</i>)	N – 46
<i>denudata dzhalilovi</i> Moravec et Amin, 1978 (<i>Rhabdochona</i>)	N – 40
<i>denudata honshuensis</i> Moravec et Nagasawa, 1989 (<i>Rhabdochona</i>)	N – 358

<i>depressa</i> (Siebold, 1836) (<i>Liga</i>)	C – 149
<i>depressum</i> (Zeder, 1800) (<i>Porrocaecum</i>)	N – 396
<i>derjavini</i> Mikailov, 1975 (<i>Gyrodactylus</i>)	M – 145
<i>dessetae</i> Baruš et Tenora, 1976 (<i>Thubunaea</i>)	N – 541
<i>diacantha</i> Petter, 1961 (<i>Syncuaria</i>)	N – 444
<i>diaphana</i> (Cholodkowski, 1906) (<i>Soricina</i>)	C – 116
<i>didelphidis</i> Caneda-Guzmán de Chambrier et Scholz, 2001 (<i>Thaumasioscolex</i>)	C – 331
<i>didelphis</i> (Butterworth et Beverley-Burton, 1977) (<i>Eucoleus</i>)	N – 257
Didymozooidea gen. sp. (<i>Torticaecum</i>) larv.	D – 79
<i>dietzevi</i> (Isaitschikov, 1927) (<i>Echinochasmus</i>)	D – 569
<i>difformis</i> Wagener, 1857 (<i>Dactylogyrus</i>)	M – 31
<i>diformoides</i> Gläser et Gussev, 1971 (<i>Dactylogyrus</i>)	M – 13
<i>digitata</i> (Linstow, 1906) (<i>Setaria</i>)	N – 476
<i>digitatus</i> (Linstow, 1906) (<i>Mecistocirrus</i>)	N – 378
<i>diminuta</i> Stunkard et Haviland, 1924 (<i>Ascocotyle</i>)	D – 342
<i>diminuta</i> Stunkard et Haviland, 1924 (<i>Ascocotyle</i>) mtc.	D – 343
<i>diminuta</i> (Rudolphi, 1819) Weinland, 1858 (<i>Hymenolepis</i>)	C – 200
<i>diodontis</i> Yamaguti, 1942 (<i>Sclerodistomum</i>)	D – 10
<i>diploorchis</i> (Yamaguti, 1936) (<i>Palaeorchis</i>)	D – 455
<i>dirigerus</i> Gussev, 1966 (<i>Dactylogyrus</i>)	M – 95
<i>discoidea</i> (Beneden, 1868) (<i>Anomotaenia</i>)	C – 68
<i>dispar</i> (Goeze, 1782) (<i>Nematotaenia</i>)	C – 21
<i>dispar</i> (Dujardin, 1845) (<i>Eucoleus</i>)	N – 321
<i>dispar</i> (Schrank, 1790) (<i>Gonguleterakis</i>)	N – 337
<i>dispar</i> (Mueller, 1936) (<i>Urocleidus</i>)	M – 264
<i>distinguendus</i> Nybelin, 1937 (<i>Dactylogyrus</i>)	M – 20
<i>distortus</i> Fusco et Overstreet, 1978 (<i>Ascarophis</i>)	N – 6
<i>ditremum</i> (Creplin, 1825) (<i>Diphyllobothrium</i>) larv.	C – 347
<i>dogieli</i> (Koval, 1950) (<i>Allocreadium</i>)	D – 120
<i>dogieli</i> Krotas, 1959 (<i>Cucullanus</i>)	N – 67
<i>dogieli</i> Pigulewsky, 1953 (<i>Phyllodistomum</i>)	D – 55
<i>donicum</i> Issaitschikoff 1909 (<i>Brachylecithum</i>)	D – 620
<i>donicus</i> Skrjabin et Lindrop, 1919 (<i>Apophallus</i>)	D – 559
<i>dorsolineatum</i> (Reimer, 1985) (<i>Steringophorus</i>)	D – 540
<i>dubia</i> (Johnston et Mawson, 1940) (<i>Cardiofilaria</i>)	N – 839
<i>dubininae</i> Matevossian, 1969 (<i>Multiuterina</i>)	C – 161
<i>dubini</i> Vojtek et Vojtková, 1968 (<i>Holostephanus</i>)	D – 296
<i>dugesii</i> Sanchez Gumiel, Zapatero Ramos, Castaño Fernandez, Gonzalez Santiago, 1991 (<i>Spinicauda</i>)	N – 531
<i>dujardini</i> (Maupas in Seurat, 1916) (<i>Entomelas</i>)	N – 95
<i>dujardini</i> (Krabbe, 1869) (<i>Aploparaksis</i>)	C – 74
<i>dukae</i> (Holl, 1928) (<i>Cosmocercoides</i>)	N – 139
<i>dulkeiti</i> Bychowsky, 1936 (<i>Dactylogyrus</i>)	M – 70, M - 392
<i>dulmaae</i> Ergens, 1970 (<i>Gyrodactylus</i>)	M – 210
<i>dyki</i> Ergens et Lucký, 1959 (<i>Dactylogyrus</i>)	M – 58
<i>dykovaе</i> Ergens, 1991 (<i>Gyrodactylus</i>)	M – 206
<i>dzhalilovi</i> Ergens et Ashurova, 1984 (<i>Gyrodactylus</i>)	M – 200
<i>ectorchis</i> (Fischthal et Kuntz, 1959) (<i>Basidiodiscus</i>)	D – 7
<i>editus</i> Dzhalilov et Ashurova, 1980 (<i>Gyrodactylus</i>)	M – 151

<i>echinobothrida</i> (Méglin, 1881) (<i>Raillietina</i>)	C – 180
<i>echinorhyncoïdes</i> (Sousin, 1889) (<i>Joyevxiella</i>)	C – 356
<i>elegans</i> Nordmann, 1832 (<i>Gyrodactylus</i>)	M – 148
<i>elegans</i> Bychowsky, 1957 (<i>Lamellodiscus</i>)	C – 266
<i>elegans</i> (Olsson, 1869) (<i>Multicalyx</i>)	D – 556
<i>elegans</i> Rudolphi, 1802 (<i>Plagiorchis</i>)	D – 220
<i>elegans</i> Gutierrez, 1945 (<i>Rhabdias</i>)	N – 215
<i>elegantis</i> Gussev, 1966 (<i>Dactylogyrus</i>)	M – 96
<i>elisae</i> (Skrjabin, 1914) (<i>Diorchis</i>)	C – 93
<i>elisae</i> Skrjabin, 1914 (<i>Aploparaksis</i>)	C – 75
<i>ellipticum</i> (Pratt, 1903) (<i>Ochetosoma</i>)	D – 110
<i>elongata</i> (Baird, 1858, Baker, 1986) (<i>Megalobatrachonema</i>)	N – 743
<i>elongatum</i> var. <i>kotlani</i> (Ihle, 1920) (<i>Cylicocyclus</i>)	N – 246
<i>elongatum</i> (Harrah, 1921) (<i>Morishitium</i>)	D – 523
<i>elongatum</i> Nybelin, 1926 (<i>Phyllodistomum</i>)	D – 54
<i>elongatus</i> (Gato et Ozaki, 1930) (<i>Neoplagioporus</i>)	D – 453
<i>emarginata</i> (Diesing, 1850) (<i>Scholzia</i>)	C – 414
<i>epinepheli</i> Scholz, Garippa et Scala, 1993 (<i>Grillotia</i>)	C – 231
<i>epinepheli</i> Moravec, Mendoza-Franco et Vargas-Vázquez, 1995 (<i>Paracapillaria</i>)	N – 643
<i>epinepheli</i> (Manter, 1947) (<i>Postporus</i>)	D – 367
<i>equispiculata</i> Moravec et Scholz, 1991 (<i>Rhabdochona</i>)	N – 556
<i>ergensi</i> Molnár, 1964 (<i>Dactylogyrus</i>)	M – 50
<i>ergensi</i> Euzet et Oliver, 1966 (<i>Lamellodiscus</i>)	M – 267
<i>ergensi</i> Scholz, 1990 (<i>Caryophyllaeides</i>)	C – 217
<i>ergensi</i> Moravec, 1968 (<i>Rhabdochona</i>)	N – 47
<i>erhardovae</i> Ergens, 1970 (<i>Dactylogyrus</i>)	M – 93
<i>erhardovae</i> Ryšavý, 1954 (<i>Prosthodendrium</i>)	D – 177
<i>erinacei</i> (Gmelin, 1790) (<i>Rodentolepis</i>)	C – 198
<i>erinacei</i> (Meggit, 1920) (<i>Mathevotaenia</i>)	C – 203
<i>erinacei europaei</i> (Rudolphi, 1819) (<i>Diphyllobothrium</i>)	C – 109
<i>erinaceus</i> (Beneden, 1858) (<i>Grillotia</i>)	C – 11
<i>erraticus</i> Rudolphi, 1809 (<i>Ichthyocotylurus</i>)	D – 603
<i>ersinensis</i> Spassky et Roytman, 1960 (<i>Dactylogyrus</i>)	M – 72
<i>emydis</i> (Liedy, 1851) (<i>Neoechinorhynchus</i>)	A – 43
<i>ensicaudatum</i> (Zeder, 1800) (<i>Porrocaecum</i>)	N – 116
<i>ephemera</i> (Nitzsch, 1817) (<i>Notocotylus</i>)	D – 201
<i>ephemeridarum</i> (Linstow, 1872) (<i>Cystidicoloides</i>)	N – 80
<i>europaea</i> Odening, 1963 (<i>Ophiotaenia</i>)	C – 24
<i>europaeum</i> Roman et Bychowsky, 1956 (<i>Octomacrum</i>)	M – 271
<i>eurubescens</i> Linton, 1910 (<i>Barisomum</i>)	D – 370
<i>eurycheilum</i> (Olsen, 1952) (<i>Hysterothylacium</i>)	N – 609
<i>euryoptera</i> (Rudolphi, 1819) (<i>Viguiera</i>)	N – 491
<i>euscorpioni</i> Baruš et Koubková, 2002 (<i>Thelastoma</i>)	N – 788
<i>ewingi</i> Chisholm et Whittington, 2000 (<i>Heliocotyle</i>)	M – 357
<i>exaeretus</i> Dietz, 1909 (<i>Petasiger</i>)	D – 209
<i>exasperatum</i> (Rudolphi, 1819) (<i>Rubenstrema</i>)	D – 328
<i>exasperatum</i> (Rudolphi, 1819) (<i>Neoglyphe</i>)	D – 505
<i>excavata</i> (Rudolphi, 1803) (<i>Tylodelphys</i>)	D – 476
<i>exigua</i> (Dujardin, 1845) López-Neyra, 1946 (<i>Aonchotheca</i>)	N – 470

<i>exilis</i> (Dujardin, 1845) (<i>Aonchotheca</i>)	N – 229
<i>exiguus</i> La Rue, 1911 (<i>Proteocephalus</i>)	C – 30
<i>expansa</i> (Rudolphi, 1805) (<i>Moniezia</i>)	C – 107
<i>expansus</i> (Creplin, 1842) (<i>Scaphanocephalus</i>) mtc.	D – 364
<i>extensus</i> Mueller et Van Cleave, 1932 (<i>Dactylogyrus</i>)	M – 5
<i>faba</i> (Bremser in Schmalz, 1831) (<i>Collyriclum</i>)	D – 522
<i>facicirrus</i> Lim, 1987 (<i>Cornudiscoides</i>)	M – 309
<i>facilis</i> (Meggitt, 1927) (<i>Liga</i>)	C – 148
<i>fairporti</i> Van Cleave, 1921 (<i>Gyrodactylus</i>)	M – 207
<i>falcatus</i> (Frölich, 1789) (<i>Acanthocephalus</i>)	A – 4
<i>falcatus</i> (Wedl, 1857) (<i>Dactylogyrus</i>)	M – 64
<i>falcatus</i> Onji et Nishio, 1916 (<i>Stellantchasmus</i>)	D – 255
<i>falcatus</i> Onji et Nishio, 1916 (<i>Stellantchasmus</i>) mtc.	D – 273
<i>falciformis</i> Achmerov, 1952 (<i>Dactylogyrus</i>)	M – 80
<i>falconi</i> Borgarenko, 1972 (<i>Euparadistomum</i>)	D – 535
<i>falconis</i> (Goeze, 1782) (<i>Baruscapillaria</i>)	N – 328, N – 504, N – 769
<i>falconis</i> Szidat, 1928 (<i>Strigea</i>)	D – 503
<i>fallax</i> Wagener, 1857 (<i>Dactylogyrus</i>)	M – 66
<i>fallax</i> (Rudolphi, 1819) (<i>Anisocladium</i>)	D – 242
<i>farciminoso</i> (Goeze, 1782) (<i>Variolepis</i>)	C – 192
<i>fariai</i> Travassos, 1914 (<i>Ornithostrongylus</i>)	N – 295
<i>farionis</i> Fischer, 1798 (<i>Cystidicola</i>)	N – 65
<i>farionis</i> (Müller, 1874) (<i>Crepidostomum</i>)	D – 43
<i>fasciata</i> (Rudolphi, 1819) (<i>Helicometra</i>)	D – 491
<i>fasciati</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 292
<i>fasciculata</i> (Ransom, 1909) (<i>Hymenolepis</i>)	C – 136
<i>fasciculi</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 291
<i>fasciolaris</i> (Pallas, 1781) (<i>Fimbriaria</i>)	C – 133
<i>fedtschenkoi</i> Skrjabin, 1924 (<i>Tanaisia</i>)	D – 142
<i>filiformis</i> (Stossich, 1896) (<i>Philometra</i>)	N – 808
<i>felineus</i> (Rivolta, 1884) (<i>Opisthorchis</i>)	D – 252, D – 411
<i>fennica</i> (Schneider, 1902) (<i>Caryophyllaeides</i>)	C – 1
<i>ferox</i> (Rudolphi, 1795) (<i>Chaunocephalus</i>)	D – 133
<i>filamentosa</i> Justine, 2004 (<i>Huffmanella</i>)	N – 817
<i>filicollis</i> (Rudolphi, 1802) (<i>Proteocephalus</i>)	C – 260
<i>filiformis</i> (Goeze, 1782) (<i>Oswaldocruzia</i>)	N – 96
<i>fimbriata</i> (Diesing, 1850) (<i>Duthiersia</i>)	C – 296
<i>fimbriatum</i> Essex, 1927 (<i>Corallobothrium</i>)	C – 270
<i>fimbriceps</i> Annenkova-Chlopina, 1919 (<i>Caryophyllaeus</i>)	C – 3
<i>finitimus</i> Gussev, 1966 (<i>Dactylogyrus</i>)	M – 63
<i>flagellicirrus</i> Gussev, 1955 (<i>Dactylogyrus</i>)	M – 81
<i>flexicollis</i> Kabata, 1961 (<i>Bathycreadium</i>)	D – 487
<i>flexis</i> (Dubois, 1934) (<i>Parastrigea</i>)	D – 580
<i>floridense</i> (Manter, 1934) (<i>Lecithochirium</i>)	D – 9
<i>fluviatile</i> Moravec et Sey, 1988 (<i>Hysterothylacium</i>)	N – 210
<i>fluviatilis</i> Bangham, 1925 (<i>Proteocephalus</i>)	C – 364
<i>fluviatilis</i> Akhmerov, 1960 (<i>Schyzocotyle</i>)	C – 204
<i>foliacea</i> (Rudolphi, 1819) (<i>Amphilina</i>)	C – 44
<i>foliaceus</i> (Linnaeus, 1758) (<i>Augulus</i>)	Cr - 7

<i>foliaceus</i> Lim et Furtado, 1985 (<i>Sundanonchus</i>)	M – 284
<i>folium</i> (Olfers, 1916) (<i>Phyllodistomum</i>)	D – 53
<i>folkmanovae</i> Ergens, 1956 (<i>Dactylogyrus</i>)	M – 61
<i>fontanus</i> (Lyster, 1939) (<i>Ptychogonimus</i>)	D – 26
<i>forcipata</i> Rudolphi, 1819 (<i>Subulura</i>)	N – 422
<i>formosa</i> (Dubinina, 1953) (<i>Microsomacanthus</i>)	C – 154
<i>formosanus</i> (Nishigori, 1924) (<i>Centrocestus</i>) mtc.	D – 271
<i>formosanus</i> (Nishigori, 1924) (<i>Centrocestus</i>)	D – 419
<i>formosus</i> Kulwiec, 1927 (<i>Dactylogyrus</i>)	M – 39
<i>fortunatowi</i> Dinnik, 1933 (<i>Rhabdochona</i>)	N – 34
<i>fossilis</i> Lupu et Roman, 1956 (<i>Gyrodactylus</i>)	M – 138
<i>fossor</i> Jägerskiöld, 1902 (<i>Dycheline</i>)	N – 19
<i>fragile</i> (Rudolphi, 1802) (<i>Eubothrium</i>)	C – 398
<i>fragilis</i> Coil, Reid et Kuntz, 1965 (<i>Paucivitellosus</i>)	D – 378
<i>fratermus</i> Wagener, 1910 (<i>Dactylogyrus</i>)	M – 14
<i>fraterna</i> Stiles, 1906 (<i>Hymenolepis</i>)	C – 392
<i>fraterna</i> (Stiles, 1906) (<i>Rodentolepis</i>)	C – 197
<i>freemani</i> (Moravec, Margolis et McDonald, 1981) (<i>Piscicapillaria</i>)	N – 1
<i>fringilae</i> Layman, 1923 (<i>Lyperosomum</i>)	D – 171
<i>fritschi</i> (Travassos, 1914) (<i>Capillostrongyloides</i>)	N – 52
<i>frontina</i> (Dujardin, 1845) (<i>Raillietina</i>)	C – 178
<i>frugilegi</i> (Czaplinski, 1962) (<i>Eucoleus</i>)	N – 322
<i>fulicae</i> (Rudolphi, 1819) (<i>Amidostomum</i>)	N – 230
<i>fulvidraconis</i> Li, 1935 (<i>Procamallanus</i>)	N – 199
<i>furcatus</i> Van Cleave et Lincicome, 1939 (<i>Illiosentis</i>)	A – 61
<i>furcatus</i> (Bremser in Rudolphi, 1819) (<i>Opecoeloides</i>)	D – 492
<i>furciger</i> (Olsson, 1868) (<i>Steringophorus</i>)	D – 493
<i>furcigera</i> (Nitzsch in Rudolphi, 1819) (<i>Aploparaksis</i>)	C – 76
<i>fuscatum</i> (Rudolphi, 1819) (<i>Brachylaima</i>)	D – 445
<i>fuscovenosa</i> (Railliet, 1889) (<i>Rhabdias</i>)	N – 94
<i>gabonica</i> Beddard, 1913 (<i>Ophiotaenia</i>)	C – 23
<i>gadi</i> Müller, 1776 (<i>Echinorhynchus</i>)	A – 7
<i>galeata</i> (Rudolphi, 1819) (<i>Rhipidocotyle</i>)	D – 489
<i>galli</i> (Schrank, 1788) (<i>Ascaridia</i>)	N – 406
<i>gallinarum</i> (Schrank, 1788) (<i>Heterakis</i>)	N – 227
<i>gallinum</i> Witenberg, 1923 (<i>Postharmostomum</i>)	D – 168
<i>garridoi</i> Baruš, 1968 (<i>Victorocara</i>)	N – 290
<i>gasterostei</i> Gläser, 1974 (<i>Gyrodactylus</i>)	M – 150
<i>gavium</i> (Guberlet, 1922) (<i>Diplostomum</i>)	D – 178
<i>geckonum</i> (Tubangui, 1929) (<i>Paradistomoides</i>)	D – 280
<i>gemini</i> Luus-Powell, Mashego et Khalil, 1994 (<i>Mormyrogyrodactylus</i>)	M – 348
<i>gibbosus</i> (Rudolphi, 1802) (<i>Lecithaster</i>)	D – 486
<i>gibbus</i> (Mehlis et Creplin, 1846) (<i>Notocotylus</i>)	D – 169
<i>gibsoni</i> De et Maity, 1994 (<i>Indocapillaria</i>)	N – 623
<i>gibsoni</i> Rego et Pavanelli, 1991 (<i>Proteocephalus</i>)	C – 407
<i>giganteum</i> (Essex, 1928) (<i>Megathylacoides</i>)	C – 26
<i>gigantica</i> (Cobbold, 1855) (<i>Fasciola</i>)	D – 251
<i>gigi</i> Fujita, 1928 (<i>Spinitectus</i>)	N – 813
<i>gemina</i> Font Overstreet et Heard, 1984 (<i>Ascocotyle</i>)	D – 448

<i>genata</i> Looss, 1907 (<i>Pygidiopsis</i>)	D – 615
<i>gerreroensis</i> Caspeta-Mandujano, Aguilaz-Aguilaz et Salgado-Maldonado, 2003 (<i>Rhabdochona</i>)	N – 793
<i>giranensis</i> (Sugimoto, 1934) (<i>Retinometra</i>)	C – 98
<i>girardi</i> Lane, 1917 (<i>Gireterakis</i>)	N – 367
<i>glaeseri</i> Ergens, 1980 (<i>Gyrodactylus</i>)	M – 256
<i>glehnii</i> Ergens et Yukhimenko, 1973 (<i>Gyrodactylus</i>)	M – 137
<i>globata</i> (Linstow, 1879) (<i>Anonchotaenia</i>)	C – 61
<i>globiceps</i> Yamaguti, 1935 (<i>Anguillicola</i>) larv.	N – 698
<i>globiceps</i> (Rudolphi, 1819) (<i>Philometra</i>)	N – 460
<i>globifera</i> (Batsch, 1786) (<i>Cladotaenia</i>)	C – 82
<i>globiporum</i> (Rudolphi, 1802) (<i>Sphaerostoma</i>)	D – 512
<i>globipunctata</i> (Rivolta, 1874) (<i>Stilesia</i>)	C – 227
<i>globocephalus</i> Mackin, 1927 (<i>Dracunculus</i>)	N – 255
<i>globuliforme</i> Cooper, 1914 (<i>Haplobothrium</i>)	C – 223
<i>globulus</i> (Rudolphi, 1819) (<i>Sphaeridiotrema</i>)	D – 285
<i>glycirrhiza</i> Beneden, 1873 (<i>Strongylacantha</i>)	N – 412
<i>glyptosterni</i> (Moravec et Amin, 1978) (<i>Alloglossidium</i>)	D – 52
<i>gnathopogonis</i> Yamaguti, 1963 (<i>Dactylogyrus</i>)	M – 83
<i>gnedini</i> Skrjabin, 1946 (<i>Rhabdochona</i>)	N – 33
<i>gobiensis</i> Gläser, 1974 (<i>Gyrodactylus</i>)	M – 229
<i>gobii</i> Schulman, 1953 (<i>Gyrodactylus</i>)	M – 146
<i>gobiomori</i> Moravec, Salgado-Maldonado et Caspeta-Mandujano, 1999 (<i>Procamallanus</i>)	N – 72
<i>gobiorum</i> Dogiel et Bychowsky, 1939 (<i>Proteocephalus</i>)	C – 299
<i>gongyla</i> (Cohn, 1900) (<i>Paricterotaenia</i>)	C – 172
<i>gonzalezi</i> Simón-Vicente, Mas-Coma, López-Roman, Tenora et Gállego, 1985 (<i>Notocotylus</i>)	D – 192
<i>golvani</i> Salgado-Maldonado, 1978 (<i>Neoechinorhynchus</i>)	A – 52
<i>goppo</i> Ozaki, 1925 (<i>Genarchopsis</i>)	D – 265
<i>gordoni</i> Codero del Campillo et Alvarez-Pelitero, 1976 (<i>Spinitectus</i>)	N – 135
<i>gotoi</i> (Motomura, 1927) (<i>Paracaryophyllaeus</i>)	C – 218
<i>gracilescens</i> (Rudolphi, 1819) (<i>Prosorhynchoides</i>)	D – 490
<i>gracilis</i> (Bellingham, 1840) (<i>Capillaria</i>)	N – 370
<i>gracilis</i> (Zeder, 1803) (<i>Sobolevicanthus</i>)	C – 183
<i>gracilis</i> (Rudolphi, 1819) (<i>Apatemon</i>)	D – 99
<i>gracillimus</i> (Lühe, 1909) (<i>Metaleptophallus</i>)	D – 76
<i>grandiacetabularis</i> Moravec et Sey, 1989 (<i>Dolichosaccus</i>)	D – 233
<i>grandis</i> Baschkirova, 1946 (<i>Echinostoma</i>)	D – 183
<i>grandivesicularis</i> Ishii, 1935 (<i>Petasiger</i>)	D – 619
<i>granulosum</i> Looss, 1907 (<i>Lecithodendrium</i>)	D – 179
<i>granulosus</i> (Batsch, 1786) (<i>Echinococcus</i>)	C – 228
<i>grayi</i> (Verma, 1936) (<i>Posthodiplostomum</i>) mtc.	D – 274
<i>groschafti</i> Ergens, 1973 (<i>Gyrodactylus</i>)	M – 136
<i>groschafti</i> Moravec, 2001 (<i>Pseudoneodiplostomum</i>)	D – 441
<i>guantanamera</i> Spiridonov, 1988 (<i>Carnoya</i>)	N – 226
<i>gueyeii</i> Koubková, Baruš et Koubek, 2005 (<i>Thelastoma</i>)	N – 841
<i>gundlachi</i> Viguera, 1943 (<i>Monoecocestus</i>)	C – 112
<i>gurleyi</i> Price, 1937 (<i>Gyrodactylus</i>)	M – 313

<i>gussevi</i> Ling Mo-en, 1962 (<i>Gyrodactylus</i>)	M – 334
<i>guttatusi</i> Andrade-Salas, Pineda-López et García-Magaña, 1994 (<i>Procamallanus</i>)	N – 825
<i>gvozdevi</i> Ergens et Kartunova (<i>Gyrodactylus</i>)	M – 341
<i>gvozdevi</i> Maksimova, 1988 (<i>Wardium</i>)	C – 210
<i>gymnosardae</i> Moravec, 2006 (<i>Philometra</i>)	N – 865
<i>haasi</i> Witenberg, 1944 (<i>Transversotrema</i>)	D – 379
<i>habanensis</i> Freze et Ryšavý, 1976 (<i>Ophiotaenia</i>)	C – 225
<i>habanensis</i> (<i>Batracholandros</i>)	N – 382
<i>haiteji</i> Ergens et Yukhimenko, 1986 (<i>Gyrodactylus</i>)	M – 168
<i>hakubaensis</i> Shimazu, 1999 (<i>Metagonimus</i>)	D – 416
<i>hakyi</i> Moravec et Sey, 1988 (<i>Rhabdochona</i>)	N – 203
<i>hamatospicula</i> Walton, 1940 (<i>Aplectana</i>)	N – 212
<i>hampalae</i> Moravec et Scholz, 1991 (<i>Camallanus</i>)	N – 552
<i>hamulatum</i> (Moulton, 1931) (<i>Cottocomephoronema</i>)	N – 56
<i>haplogonoides</i> Gussev, 1966 (<i>Dactylogyrus</i>)	M – 65
<i>harpiprioni</i> Georgiev et Vaucher, 2004 (<i>Arnelepis</i>)	C – 379
<i>hartwichi</i> Moravec, Wolter et Körting, 1999 (<i>Dichelyne</i>)	N – 732
<i>hawaiiense</i> (Yamaguti, 1970) (<i>Weketrema</i>)	D – 451
<i>hayesi</i> (<i>Didelphostrongylus</i>)	N – 433
<i>hebraicus</i> Dubois, 1934 (<i>Cotylurus</i>)	D – 533
<i>helicina</i> (Molin, 1860) (<i>Dujardinascaris</i>)	N – 238
<i>helicoleni</i> Bray et Kuchta 2006 (<i>Brachyenteron</i>)	D – 558
<i>hellichi</i> (Šrámek, 1901) (<i>Rhabdochona</i>)	N – 48
<i>hellichi</i> (Šrámek, 1901) (<i>Rhabdochona</i>) – larvae from <i>Hydropsyche</i>	N – 682
<i>helocirrus</i> Barton, 1994 (<i>Dolichosaccus</i>)	D – 305
<i>hemiamphibothrium</i> Ergens, 1956 (<i>Dactylogyrus</i>)	M – 59
<i>hemibarbi</i> Ergens, 1980 (<i>Gyrodactylus</i>)	M – 333
<i>hemiculteris</i> Ergens et Gussev, 1975 (<i>Gyrodactylus</i>)	M – 129
<i>hemidactyli</i> Moravec et Baruš, 1990 (<i>Skrjabinodon</i>)	N – 459
<i>appendiculatus</i> (Rudolphi, 1802) (<i>Hemiurus</i>)	D – 495
<i>hemivicinus</i> Ergens et Daniyarov, 1976 (<i>Gyrodactylus</i>)	M – 228
<i>hepatica</i> Linnaeus, 1758 (<i>Fasciola</i>)	D – 151
<i>hermaphrodita</i> (Froel, 1789) (<i>Ascaridia</i>)	N – 508
<i>herodiae</i> Fuhrmann, 1912 (<i>Dendrouterina</i>)	C – 417
<i>herotilapiae</i> Vidal-Martínez, Scholz et Aguirre-Macedo, 2001 (<i>Gussevia</i>)	M – 362
<i>heteranchorus</i> (Kulkarni, 1969) (<i>Mastacembelocleidus</i>)	M – 404
<i>heterocerca</i> (Goto, 1894) (<i>Heteraxine</i>)	M – 407
<i>heterolecithodes</i> (Braun, 1899) (<i>Athesmia</i>)	D – 170
<i>heterostomum</i> Rudolphi, 1809 (<i>Euclinostomum</i>)	D – 532
<i>heuttoneni</i> Haukisalml et Tenora, 1993 (<i>Catenotaenia</i>)	C – 236
<i>hians</i> (Rudolphi, 1819) (<i>Cathaemasia</i>)	D – 534
<i>higai</i> Yamaguti, 1936 (<i>Philopinna</i>)	D – 480
<i>hilarii</i> Vaz et Pereira, 1934 (<i>Procamallanus</i>)	N – 607
<i>himantopodis</i> (Krabbe, 1869) (<i>Hymenolepis</i>)	C – 137
<i>himezi</i> Yamaguti, 1941 (<i>Cucullanus</i>)	N – 253
<i>hippopotami</i> Stunkard, 1924 (<i>Oculotrema</i>)	M – 396
<i>hirsutum</i> (Looss, 1896) (<i>Lecithodendrium</i>)	D – 101
<i>hirudinaceus</i> (Pallas, 1781) (<i>Macracathorhynchus</i>)	A – 41
<i>hirundina</i> (Fuhrmann, 1907) (<i>Anomotaenia</i>)	C – 69

<i>hispida</i> (Van Cleave, 1925) (<i>Southwellina</i>)	A – 50
<i>hispida</i> (Van Cleave, 1925) (<i>Southwellina</i>) larv.	A – 58
<i>hispidum</i> Fedtschenko, 1872 (<i>Gnathostoma</i>)	N – 104
<i>hoeplii</i> Hsü, 1935 (<i>Oswaldocruzia</i>)	N – 164
<i>hokkaidensis</i> Asakawa, Kamiya et Obayashi, 1988 (<i>Zonorchis</i>)	D – 239
<i>hokkaidensis</i> Asakawa, Kamiya et Obayashi, 1988 (<i>Liniscus</i>)	N – 275
<i>hollisae</i> Groschaft, 1972 (<i>Bravotrema</i>)	D – 154
<i>holopterum</i> (Rudolphi, 1819) (<i>Angusticaecum</i>)	N – 600
<i>holostomum</i> (Rudolphi, 1819) (<i>Leucochloridium</i>)	D – 182
<i>homoin gracile</i> (Reichenbach-Klinke, 1961) (<i>Paradiplozoon</i>)	M – 305
<i>homoin homoin</i> (Bychowsky et Nagibina, 1959) (<i>Paradiplozoon</i>)	M – 306
<i>honshuensis</i> Moravec et Nagasawa, 1998 (<i>Plagioporus</i>)	D – 406
<i>horrida</i> Rudolphi, 1809 (<i>Skrjabinoclava</i>)	N – 294
<i>horrida</i> (Linstow, 1901) (<i>Hymenolepis</i>)	C – 113
<i>hospeti</i> Thapar, 1950 (<i>Rhabdochona</i>)	N – 39
<i>howelli</i> (Viguera, 1934) (<i>Pseudohelimosomum</i>)	N – 418
<i>hrabei</i> Ergens, 1957 (<i>Gyrodactylus</i>)	M – 174
<i>huffmani</i> Moravec, 1986 (<i>Huffmanella</i>)	N – 157
<i>huffmani</i> Moravec, 2000 (<i>Paracapillaria</i>)	N – 750
<i>huchonis</i> Bauer, 1948 (<i>Tetraonchus</i>)	M – 270
<i>humbertoi</i> Caspeta-Mandujano et Moravec, 2000 (<i>Spinitectus</i>)	N – 766
<i>humili</i> Roytman et Trofimenko, 1964 (<i>Rhabdochona</i>)	N – 42
<i>huronensis</i> Uhazy, 1976 (<i>Philometroides</i>)	N – 28
<i>hwangtsiyui</i> Tsing, 1933 (<i>Azygia</i>)	D – 212
<i>hydatigena</i> Pallas, 1776 (<i>Taenia</i>)	C – 115
<i>hypoprioni</i> Potter, 1927 (<i>Platybothrium</i>)	C – 323
<i>hypostomi</i> Petter et Cassone, 1984 (<i>Raphidascaris</i>)	N – 458
<i>hypophthalmichthys</i> Dogiel et Akhmerov, 1959 (<i>Camallanus</i>)	N – 810
<i>hypophthalmichthys</i> Achmerow, 1952 (<i>Dactylogyrus</i>)	M – 68
<i>hypophthalmichthysi</i> Ling, 1962 (<i>Gyrodactylus</i>)	M – 235
<i>hypselobagri</i> (Billet, 1898) (<i>Isoparorchis</i>)	D – 213
<i>ibisae</i> (Schmidt et Bush, 1972) (<i>Cyclusteria</i>)	C – 280
<i>ibisae</i> (Schmidt et Bush, 1972) (<i>Cyclusteria</i>) larv.	C – 337
<i>ibragimovi</i> Ergens, 1980 (<i>Gyrodactylus</i>)	M – 255
<i>icosiensis</i> (Seurat, 1917) (<i>Moaciria</i>)	N – 170
<i>iheringi</i> (Travassos, Artigas et Pereira, 1928) (<i>Klossinemella</i>)	N – 679
<i>iliensis</i> Gvosdev et Martechov, 1953 (<i>Paragyrodactylus</i>)	M – 281
<i>illensis</i> (Ziegler, 1883) (<i>Rhipidocotyle</i>)	D – 2
<i>illiciens</i> (Braun, 1901) (<i>Platynosomum</i>)	D – 156
<i>imbricatus</i> (Looss, 1893) (<i>Notocotylus</i>)	D – 202
<i>imias</i> Spiridonov, 1988 (<i>Heth</i>)	N – 264
<i>imitans</i> (Mühling, 1898) (<i>Asymphyrodora</i>)	D – 48
<i>immitis</i> (Leidy, 1856) (<i>Dirofilaria</i>)	N – 477
<i>immutatum</i> Ward et Magath, 1917 (<i>Haplonema</i>)	N – 254
<i>impar</i> Bray, Cribb (<i>Preptetos</i>)	D – 346
<i>inamata</i> de Chambrier, Al-Kallak et Mariáns, 2003 (<i>Postgangesia</i>)	C – 422
<i>incerta</i> Caballero, 1949 (<i>Aplectana</i>)	N – 729
<i>incognita</i> Solonitsin, 1932 (<i>Desmidocercella</i>)	N – 286
<i>incognitus</i> Szidat, 1943 (<i>Palaeorchis</i>)	D – 69

<i>incognitus</i> Ergens et Gussev, 1980 (<i>Gyrodactylus</i>)	M – 253
<i>incurvum</i> (Rudolphi, 1819) (<i>Hysterothylacium</i>)	N – 148
<i>indica</i> Moravec, Arthur, Razia Beevi et Radhakrishnan, 1993 (<i>Pseudocapillaria</i>)	N – 597
<i>inermicauda</i> Baylis, 1923 (<i>Pharyngodon</i>)	N – 181
<i>inermis</i> Plehn, 1905 (<i>Sanguinicola</i>)	D – 109
<i>inermis</i> (Zeder, 1800) (<i>Spinitectus</i>)	N – 58
<i>inexpectatus</i> Izjumova et Gussev, 1955 (<i>Dactylogyrus</i>)	M – 70
<i>inflata</i> (Molin, 1859) (<i>Deropristis</i>)	D – 90
<i>inflata</i> (Rudolphi, 1819) (<i>Diorchis</i>)	C – 120
<i>inflexus</i> (Rudolphi, 1809) (<i>Pseudalius</i>)	N – 790
<i>infundibulovagina</i> Yamaguti, 1942 (<i>Ancylo-discoides</i>)	M – 110
<i>infundibulum</i> (Bloch, 1779) (<i>Choanotaenia</i>)	C – 145
<i>inimici</i> Yamaguti, 1941 (<i>Philometra</i>)	N – 702
<i>inopinatus</i> Travassos, Artigas et Pereira, 1928 (<i>Procamallanus</i>)	N – 529
<i>inopinatus</i> (cf.) (<i>Procamallanus</i>)	N – 674
<i>insigne</i> (Boulenger, 1917) (<i>Cylicocyclus</i>)	N – 247
<i>insigne</i> (Shuler, 1938) (<i>Otobothrium</i>)	C – 272
<i>insignis</i> Loeungberg, 1890 (<i>Ophryocotyle</i>)	C – 164
<i>integerimum</i> (Frölich, 1791) (<i>Polystoma</i>)	M – 2
<i>intermedia</i> (Fuhrmann, 1913) (<i>Fimbriarioides</i>)	C – 99
<i>intermedia</i> Rudolphi, 1810 (<i>Taenia</i>)	C – 199
<i>intermedius</i> Wagener, 1910 (<i>Dactylogyrus</i>)	M – 9
<i>interrupta</i> (Rudolphi, 1810) (<i>Digramma</i>)	C – 376
<i>intestinalis</i> (Dogiel et Bychowsky, 1934) (<i>Molnaria</i>)	N – 61
<i>intestinalis</i> (Linnaeus, 1758) (<i>Ligula</i>)	C – 150
<i>intestinalis</i> (Linnaeus, 1758) (<i>Ligula</i>) larv.	C – 38
<i>isabellae</i> (Lamothe-Argumedo, 1977) (<i>Genarchella</i>)	D – 311
<i>iskanderensis</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 294
<i>isoporum</i> (Looss, 1894) (<i>Allocreadium</i>)	D – 40
<i>italica</i> (Ricci, 1949) (<i>Aonchotheca</i>)	N – 324
<i>itzocanensis</i> Bravo-Hollis, 1943 (<i>Aplectana</i>)	N – 730
<i>ivanovici</i> Ergens, 1970 (<i>Dactylogyrus</i>)	M – 10
<i>izjumovae</i> Gussev, 1966 (<i>Dactylogyrus</i>)	M – 11
<i>jacksoni</i> (Bullock, 1962) (<i>Acanthocephalus</i>)	A – 3
<i>jacksonii</i> (Cobbold, 1869) (<i>Fasciola</i>)	D – 126
<i>jaliscensis</i> Moravec, Salgado-Maldonado et Caspeta-Mandujano, 1999 (<i>Procamallanus</i>)	N – 727
<i>japonensis</i> (Yamaguti, 1934) (<i>Khawia</i>)	C – 348
<i>japonensis</i> Yamaguti, 1935 (<i>Spironoura</i>)	N – 523
<i>japonica</i> Moravec, Koudela, Ogawa et Nagasawa, 1998 (<i>Huffmanella</i>)	N – 693
<i>japonica</i> (Wilkie, 1930) (<i>Meterakis</i>)	N – 208
<i>japonica</i> Yamaguti, 1934 (<i>Anguilla</i>)	C – 374
<i>japonica</i> Yamaguti, 1938 (<i>Cosmocerca</i>)	N – 163
<i>japonica</i> Moravec et Nagasawa, 1985 (<i>Ichthyofilaria</i>)	N – 154
<i>japonica</i> Moravec, 1975 (<i>Rhabdochona</i>)	N – 710
<i>japonicum</i> Moravec, Ogawa et Nagasawa, 2000 (<i>Dichelyne</i>)	N – 761
<i>japonicus</i> Kichuchi, 1929 (<i>Gyrodactylus</i>)	M – 189
<i>jiangxiensis</i> Wang, Zhao, Wang et Zhang, 1979 (<i>Rhabdochona</i>)	N – 202

<i>jimenezi</i> Ryšavý, 1966 (<i>Paraprogynotaenia</i>)	C – 170
<i>jiroveci</i> Ergens et Bychowsky, 1967 (<i>Gyrodactylus</i>)	M – 162
<i>joanae</i> de Chambrier et Paulina, 1997 (<i>Proteocephalus</i>)	C – 259
<i>joaquiensis</i> Ingles, 1936 (<i>Rhabdias</i>)	N – 138
<i>justinei</i> Moravec, Ternengo et Levron, 2006 (<i>Philometra</i>)	N – 853
<i>juvenilis</i> (Nicoll, 1918) (<i>Dolichosaccus</i>)	D – 304
<i>kafirniganensis</i> Ergens et Daniyarov, 1976 (<i>Gyrodactylus</i>)	M – 227
<i>kajika</i> Urabe et Higa, 2006 (<i>Neoplagioporus</i>)	D – 591
<i>akea</i> Bhalerao, 1926 (<i>Brachylecithum</i>)	D – 527
<i>karatagensis</i> Ergens et Allamuratov (<i>Gyrodactylus</i>)	M – 133
<i>kartasi</i> Neifar, Euzet et Hassine, 1999 (<i>Helicotyle</i>)	M – 351
<i>karvei</i> Bhalerao, 1937 (<i>Bucephalopsis</i>)	D – 266
<i>katharineri</i> Malmberg, 1964 (<i>Gyrodactylus</i>)	M – 7
<i>kaverii</i> Karve et Naik, 1951 (<i>Spironoura</i>)	N – 206
<i>kearni</i> Ergens, 1990 (<i>Gyrodactylus</i>)	M – 249
<i>kherulensis</i> Ergens, 1974 (<i>Gyrodactylus</i>)	M – 140
<i>kidderi</i> Pearse, 1936 (<i>Rhabdochona</i>)	N – 629
<i>kidderi texensis</i> Moravec et Huffman, 1988 (<i>Rhabdochona</i>)	N – 281
<i>kindderi texensis</i> (<i>Rhabdochona</i>) larv.	N – 753
<i>kisutchi</i> Margolis, Moravec et McDonald, 1975 (<i>Rhabdochona</i>)	N – 5
<i>kobayashii</i> Hukuda, 1940 (<i>Gyrodactylus</i>)	M – 143
<i>kobuleji</i> Molnár et Fernando, 1975 (<i>Philometra</i>)	N – 29
<i>koiae</i> Gibson et Bray, 1984 (<i>Anomalotrema</i>)	D – 488
<i>kolimensis</i> (Gubanov et Fedorov, 1967) (<i>Pterygodermatites</i>)	N – 438
<i>konovalovi</i> Ergens, 1976 (<i>Gyrodactylus</i>)	M – 219
<i>koreanum</i> Park, 1938 (<i>Carassotrema</i>)	D – 214
<i>koschewnikovi</i> Skryabin et Massino, 1925 (<i>Stromitrema</i>)	D – 521
<i>kotlani</i> (Molnár, 1969) (<i>Philometra</i>)	N – 73
<i>krabbeella</i> Hughes, 1940 (<i>Sobolevicanthus</i>)	C – 184
<i>krameri</i> (Petter, 1974) (<i>Procamallanus</i>)	N – 680
<i>kritscheri</i> Moravec, Salgado Maldonado et Pineda López, 1993 (<i>Raillietnema</i>)	N – 596
<i>kubanicum</i> (Issaitschikoff, 1923) (<i>Asymphylogora</i>)	D – 117
<i>kulasirii</i> (Yeh, 1960) (<i>Camallanus</i>)	N – 394
<i>kuntzi</i> Moravec et Gibson, 1986 (<i>Paracapillaria</i>)	N – 159
<i>kutassi</i> (Schulz, 1927) (<i>Streptopharagus</i>)	N – 366
<i>kutori</i> (Rukhlyadeva, 1946) (<i>Baruscapillaria</i>)	N – 243
<i>laciniatum</i> Linton, 1890 (<i>Anthobothrium</i>)	C – 320
<i>lacteum</i> (Jägerskiöld, 1896) (<i>Galactosomum</i>)	D – 298
<i>lacustris</i> Thurston et Paperna, 1969 (<i>Diplectanum</i>)	M – 260
<i>lacustris</i> (Zoega, 1776) (<i>Camallanus</i>)	N – 82
<i>laeviconchus</i> (Wedl, 1862) (<i>Procamallanus</i>)	N – 13
<i>laevis</i> (Müller, 1776) (<i>Pomphorhynchus</i>)	A – 26
<i>laevis</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 154
<i>laevis</i> (Bloch, 1782) (<i>Diploposthe</i>)	C – 129
<i>lagowskii</i> Ergens, 1980 (<i>Gyrodactylus</i>)	M – 258
<i>lamberti</i> Ergens, 1973 (<i>Gyrodactylus</i>)	M – 135
<i>lamellatus</i> Achmerow, 1952 (<i>Dactylogyrus</i>)	M – 12
<i>lamothei</i> (García-Prieto, 1990) (<i>Megathylacoides</i>)	C – 304

<i>lanceolata</i> (Bloch, 1782) (<i>Drepanidotaenia</i>)	C – 166
<i>lari</i> Blanchard, 1849 (<i>Cyathostoma</i>)	N – 776
<i>lari</i> Timon-David, 1933 (<i>Renicola</i>)	D – 143
<i>laricola</i> Skrjabin, 1924 (<i>Plagiorchis</i>)	D – 219
<i>laruei</i> Lamont, 1921 (<i>Glaridacris</i>)	C – 322
<i>laryngeus</i> Railliet, 1899 (<i>Mammomonogamus</i>)	N – 302
<i>lateolabracis</i> (Yamagiti, 1935) (<i>Philometra</i>)	N – 252
<i>laterifilamenta</i> Moravec, Kohn et Fernandes, 1992 (<i>Ichthyouris</i>)	N – 581
<i>laticeps</i> (Pallas, 1781) (<i>Caryophyllaeus</i>)	C – 2
<i>latum</i> (Linnaeus, 1758) (<i>Diphyllobothrium</i>)	C – 428
<i>latus</i> Bychowsky, 1933 (<i>Gyrodactylus</i>)	M – 160
<i>latus</i> Řehulková et Gelnar, 2005 (<i>Thylacicleidus</i>)	M – 398
<i>lavareti</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 254
<i>legendrei</i> Pariselle et Euzet, 2003 (<i>Cichlidogyrus</i>)	M – 387
<i>lemoallei</i> Pariselle et Euzet, 2003 (<i>Cichlidogyrus</i>)	M – 388
<i>388eopoldinae</i> (<i>Echinochasmus</i>)	D – 317
<i>leopoldinae</i> Scholz, Ditrich et Vargas-Vázquez, 1996 (<i>Echinochasmus</i>) mtc.	D – 331
<i>lenha</i> (Woodland, 1933) (<i>Peltidocotyle</i>)	C – 411
<i>lenoki</i> Gussev, 1953 (<i>Gyrodactylus</i>)	M – 131
<i>lenoki</i> Achmerow, 1952 (<i>Tetraonchus</i>)	M – 274
<i>lenteixeirai</i> Pérez Viguera, 1938 (<i>Oswaldocruzia</i>)	N – 214
<i>leonina</i> (Linstow, 1902) (<i>Toxascaris</i>)	N – 395
<i>leopoldinae</i> Scholz, Ditrich et Vargas-Vázquez, 1996 (<i>Echinochasmus</i>)	D – 317
<i>leporini</i> Petter, 1989 (<i>Dichelyne</i>)	N – 580
<i>leptogaster</i> (Leuckart, 1830) (<i>Chimaericola</i>)	M – 417
<i>leptoptera</i> (Rudolphi, 1819) (<i>Habronema</i>)	N – 497
<i>leptosoma</i> (Gervais, 1848) (<i>Abbreviata</i>)	N – 436
<i>leucisci</i> Žitňan, 1964 (<i>Gyrodactylus</i>)	M – 242
<i>leucisci</i> Ergens, 1963 (<i>Pseudochetosoma</i>)	D – 123
<i>levenseni</i> (Linton, 1900) (<i>Lepidapedon</i>)	D – 366
<i>liaohoensis</i> Gussev, 1962 (<i>Dactylogyrus</i>)	M – 85
<i>lichtenfelsi</i> Sánchez-Alvarez, García-Prieto et Pérez-Ponce de León, 1998(<i>Rhabdochona</i>)	N – 71
<i>limneus</i> Malmberg, 1964 (<i>Gyrodactylus</i>)	M – 178
<i>limnicola</i> Brugni et Viozzi, 2006 (<i>Pseudodelphis</i>)	N – 833
<i>liewi</i> du Preez et Lim, 2000 (<i>Neopolystoma</i>)	M – 355
<i>lingua</i> (Creplin, 1825) (<i>Cryptocotyle</i>)	D – 195
<i>linstowi</i> Dollfus, 1931 (<i>Lecithodendrium</i>)	D – 181
<i>liobagri</i> Moravec et Nagasawa, 1998 (<i>Mexiconema</i>)	N – 709
<i>litteratus</i> (Batsch, 1786) (<i>Mesocestoides</i>)	C – 397
<i>lizae</i> Lin, 2002 (<i>Lasiotocus</i>)	D – 464
<i>llewellyni</i> Ergens et Dulmaa, 1967 (<i>Gyrodactylus</i>)	M – 194
<i>lobata</i> (Fuhrmann, 1906) (<i>Skrjabinolepis</i>)	C – 111
<i>lobata</i> (Baer, 1925) (<i>Skrjabinotaenia</i>)	C - 286
<i>lobatum</i> (Wallin, 1909) (<i>Allocreadium</i>)	D – 36
<i>lobatum</i> (Railliet, 1900) (<i>Brachylecithum</i>)	D - 472
<i>lobulata</i> (Mayhew, 1925) (<i>Lobatolepis</i>)	C – 152

<i>locellus</i> Kossack, 1910 (<i>Opisthioglyphe</i>)	D – 205
<i>lomi</i> Ergens et Gelnar, 1988 (<i>Gyrodactylus</i>)	M – 332
<i>longa</i> Ranson, 1920 (<i>Ascocotyle</i>) mtc.	D – 360, D – 445
<i>longibursatum</i> (Yorke et Macfie, 1918) (<i>Trichonema</i>)	N – 543
<i>longibursatus</i> Moravec et Sey, 1989 (<i>Dolichosaccus</i>)	D – 234
<i>longicauda</i> Dzhililov, 1964 (<i>Rhabdochona</i>)	N – 32
<i>longicauda</i> (Linstow, 1885) (<i>Cosmocerca</i>)	N – 87
<i>longicaudatum</i> Brennes et Bravo Hollis, 1960 (<i>Cyrtosomum</i>)	N – 350
<i>longicaudatus</i> Moravec, 1973 (<i>Camallanus</i>)	N – 7
<i>longicaudum</i> (Rudolphi, 1809) (<i>Lyperosomum</i>)	D - 477
<i>longicirrosa</i> (Fuhrmann, 1906) (<i>Retinometra</i>)	C – 101
<i>longicollis</i> (Abildgaard, 1788) (<i>Macrodera</i>)	D – 81
<i>longicollis</i> (Rudolphi, 1819) (<i>Cardiocephalus</i>)	D – 184
<i>longicollis</i> (Zeder, 1800) (<i>Proteocephalus</i>)	C – 306
<i>longicollis</i> (Zeder, 1800) (<i>Proteocephalus</i>) larv.	C – 351
<i>longicornis</i> Paperna et Thurston, 1969 (<i>Cichlidogyrus</i>)	M – 120
<i>longifilla</i> (Dujardin, 1845) (<i>Aonchotheca</i>)	N – 774
<i>longifila</i> (Dujardin, 1845) (<i>Capillaria</i>)	N – 415
<i>longiovata</i> Moravec et Spratt, 1998 (<i>Crocodylocapillaria</i>)	N – 694
<i>longiovatus</i> Baruš, 1969 (<i>Tetrameres</i>)	N – 368
<i>longiradix</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 158
<i>longireceptaculis</i> Moravec et Sey, 1989 (<i>Paradentitruncus</i>)	A – 34
<i>longirostris</i> Moravec et Sey, 1989 (<i>Cleaveius</i>)	A – 36
<i>longispicula</i> Moravec et Kaiser, 1995 (<i>Cosmocerca</i>)	N – 601
<i>longispicula</i> (Sonsino, 1889) (<i>Paracapillaria</i>)	N – 146
<i>longispiculum</i> Moravec et Prouza, 1996 (<i>Guyanema</i>)	N – 669
<i>longissimum</i> (Ortlepp, 1923) (<i>Heliconema</i>)	N – 862
<i>longivitellaris</i> Moravec et Sey, 1989 (<i>Pleurogenes</i>)	D – 229
<i>longivitellata</i> Shtrom, 1947 (<i>Tanaisia</i>)	D – 150
<i>longleyi</i> Moravec et Huffman, 2000 (<i>Brachycoelium</i>)	D – 435
<i>longleyi</i> Moravec et Huffman, 1988 (<i>Rhabdochona</i>)	N – 280
<i>longoacuminatus</i> Žitňan, 1964 (<i>Gyrodactylus</i>)	M – 250
<i>longulum</i> Travassos, Freitas et Bührnheim, 1965 (<i>Homalometron</i>)	D – 414
<i>longus</i> (Ransom, 1920) (<i>Phagicola</i>)	D – 295
<i>longus</i> Moravec, Justine et Rigby, 2006 (<i>Procamallanus</i>)	N – 863
<i>loossi</i> (Pérez Viguera, 1955) (<i>Pelaezia</i>)	D – 308
<i>loossi</i> (Pérez Viguera, 1957) (<i>Pelaezia</i>) mtc.	D – 335
<i>lopesi</i> Rego, 1989 (<i>Nomimoscolex</i>)	C - 406
<i>lphocara</i> Gerichter, 1951 (<i>Crenosoma</i>)	N – 235
<i>lotae</i> Gussev, 1953 (<i>Gyrodactylus</i>)	M – 183
<i>lucida</i> Ransom, 1900 (<i>Metroliasthes</i>)	C – 158
<i>lucii</i> (Müller, 1776) (<i>Azygia</i>)	D – 95
<i>lucii</i> Kulakowskaja, 1951 (<i>Gyrodactylus</i>)	M – 172
<i>lucii</i> (Müller, 1776) (<i>Acanthocephalus</i>)	A – 1
<i>lucioercae</i> Gussev, 1962 (<i>Gyrodactylus</i>)	M – 155
<i>lucioercae</i> (Müller, 1776) (<i>Bunodera</i>)	D – 45
<i>lucknowensis</i> Agrawal et Sharma, 1990 (<i>Bicotyle</i>)	M – 337
<i>luckyi</i> Ergens, 1970 (<i>Gyrodactylus</i>)	M – 212
<i>lupi</i> (Rudolphi, 1819) (<i>Spirocerca</i>)	N – 187
<i>luquei</i> Kohn, Baptista-Farias et Cohen, 2000 (<i>Paranaella</i>)	M – 360

<i>luscae</i> (Beneden et Hesse, 1864) (<i>Diclidophora</i>)	M – 349
<i>macallisteri</i> Burse et Goldberg, 1996 (<i>Oochoristica</i>)	C – 251
<i>macdonaldi</i> (Dobson, 1880) (<i>Rictularia</i>)	N – 336
<i>macdonaldi</i> Rao, 1951 (<i>Paracuaria</i>)	N – 429
<i>macilenthus</i> Van Cleave, 1919 (<i>Octospinifer</i>)	A – 23
<i>macintoshii</i> (Stewart, 1914) (<i>Aplectana</i>)	N – 143
<i>maculosus</i> (Rudolphi, 1802) (<i>Plagiorchis</i>)	D – 203
<i>macracantha</i> (Joyeux et Baer, 1935) (<i>Amirthalingamia</i>)	C – 292
<i>macracanthus</i> Wagener, 1909 (<i>Dactylogyrus</i>)	M – 6
<i>macracanthus</i> Hukuda, 1940 (<i>Gyrodactylus</i>)	M – 195
<i>macroacetabularis</i> Sey, 1986 (<i>Platycladorchis</i>)	D – 498
<i>macrocaudatus</i> Ditrich, Scholz et Vargas-Vázquez, 1996 (<i>Echinochasmus</i>)	D – 318
<i>macrocephala</i> Douthitt, 1915 (<i>Andrya</i>)	C – 390
<i>macrocephalus</i> (Creplin, 1825) (<i>Proteocephalus</i>)	C – 209
<i>macrocaudatus</i> (<i>Echinochasmus</i>)	D – 318
<i>macrocaudatus</i> Ditrich, Scholz et Vargas-Vázquez, 1996 (<i>Echinochasmus</i>) mtc.	D – 332
<i>macrocolpius</i> Řehulková et Gelnar, 2006 (<i>Dactylogyrus</i>)	M – 410
<i>macrocornis</i> Ergens, 1963 (<i>Gyrodactylus</i>)	M – 187
<i>macronychus</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 163
<i>macropeos</i> (Wedl, 1855) (<i>Parvitaenia</i>)	C – 283
<i>macrophallos</i> Parona, 1889 (<i>Hastospiculum</i>)	N – 304
<i>macrophallus</i> (Diesing, 1850) (<i>Proteocephalus</i>)	C – 247
<i>macrorhodei</i> Ergens et Yukhimenko, 1975 (<i>Gyrodactylus</i>)	M – 141
<i>macrorhyncha</i> (Rudolphi, 1810) (<i>Schistotaenia</i>)	C – 182
<i>macrostoma</i> Ozaki, 1925 (<i>Asymphylogora</i>)	D – 452
<i>macrostoma</i> (Faust, 1918) (<i>Proterometra</i>)	D – 199
<i>macrostoma</i> Moravec et Mikailov, 1970 (<i>Rhabdochona</i>)	N – 108
<i>macrostomum</i> (Rudolphi, 1803) (<i>Leucochloridium</i>)	D – 146
<i>macrostomum</i> (Jägerskiöld, 1900) (<i>Pulvinifer</i>)	D – 614
<i>macrostomus</i> (Robinson, 1956) (<i>Ascocotyle</i>)	D – 350
<i>macrostomus</i> (Rudolphi, 1802) (<i>Urogonismus</i>)	D – 436
<i>macrouri</i> Moravec, Klimpel et Kara, 2006 (<i>Neoscarophis</i>)	N – 845
<i>madagascariensis</i> Chabaud, 1960 (<i>Geopetitia</i>)	N – 340
<i>magna</i> (Baer, 1959) (<i>Cyclusteria</i>) larv.	C – 293
<i>magna</i> (Bassi, 1875) (<i>Fascioloides</i>)	D – 462
<i>magniuterina</i> Ryšavý, 1957 (<i>Anonchotaenia</i>)	C – 73
<i>magnificus</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 182
<i>magnum</i> Ryšavý, 1956 (<i>Prosthodendrium</i>)	D – 172
<i>magnus</i> Konovalov, 1967 (<i>Gyrodactylus</i>)	M – 217
<i>magnus</i> (Bychowsky et Nagibina, 1957) (<i>Thaparocleidus</i>)	M – 394
<i>mahnerti</i> (Petter et Cassone, 1984) (<i>Raphidascaris</i>)	N – 461
<i>majeedi</i> Sinha et Hakim, 1967 (<i>Postorchigenes</i>)	D – 282
<i>major</i> Boudaya, Neifar et Euzet, 2006 (<i>Plectanocotyle</i>)	M – 413
<i>major</i> (Hasegawa, 1935) Yamaguti, 1938 (<i>Pseudexorchis</i>)	D – 552
<i>malayanus</i> Lim et Furtado, 1984 (<i>Dactylogyrus</i>)	M – 369
<i>malayensis</i> Lim, 1987 (<i>Cornudiscoides</i>)	M – 311
<i>malayensis</i> Lim et Furtado, 1984 (<i>Dactylogyroides</i>)	M – 374
<i>maldivensis</i> Moravec, Lorber et Konečný, 2006 (<i>Heptochona</i>)	N – 866
<i>malleus</i> Linstow, 1877 (<i>Dactylogyrus</i>)	M – 36
<i>malmbergensis</i> Prost, 1974 (<i>Gyrodactylus</i>)	M – 144

<i>malmbergi</i> Ergens, 1961 (<i>Gyrodactylus</i>)	M – 177
<i>malopteruri</i> (Fritsch, 1886) (<i>Electrotaenia</i>)	C – 373
<i>mamillana</i> (Meblis, 1831) (<i>Paranoplocephala</i>)	C – 201
<i>mamillatus</i> (Linstow, 1897) (<i>Pharyngodon</i>)	N – 180
<i>managatuwo</i> Yamaguti, 1941 (<i>Philometra</i>)	N – 783
<i>mansoni</i> (Cobbold, 1882) (<i>Diphyllobothrium</i>)	C – 110
<i>mansoni</i> Sambon, 1907 (<i>Schistosoma</i>)	D – 242
<i>mansoni</i> (Cobbold, 1879) (<i>Oxyspirura</i>)	N – 474
<i>manteri</i> Watson, 1976 (<i>Oligogonotylus</i>)	D – 315
<i>manteri</i> Watson, 1976 (<i>Oligogonotylus</i>) mtc.	D – 323
<i>mantschuricus</i> Ergens et Yukhimenko, 1977 (<i>Gyrodactylus</i>)	M – 238
<i>margolisi</i> Moravec et McDonald, 1981 (<i>Capillaria</i>)	N – 2
<i>margolisi</i> Maity et De, 1996 (<i>Discocapillaria</i>)	N – 645
<i>margolisi</i> Moravec, Vidal-Martínez et Aguirre- Macedo, 1994 (<i>Philometra</i>)	N – 622
<i>margolisi</i> Bray, 1995 (<i>Steringophorus</i>)	D – 541
<i>mariae</i> (Layman, 1930) (<i>Clavinema</i>)	N – 195
<i>marina</i> Manter, 1947 (<i>Crassicutis</i>)	D – 413
<i>maringaensis</i> Pavanelli et Rego, 1989 (<i>Spatulifer</i>)	C - 410
<i>marinum</i> (Linnaeus, 1767) (<i>Hysterothylacium</i>)	N – 739
<i>marjami</i> Allamuratov et Gussev, 1969 (<i>Gyrodactylus</i>)	M – 193
<i>markakulensis</i> Gvosdev, 1950 (<i>Gyrodactylus</i>)	M – 161
<i>markewitschi</i> Kulakowskaja, 1951 (<i>Gyrodactylus</i>)	M – 175
<i>markewitschi</i> Kulakowskaja, 1947 (<i>Asymphylogora</i>)	D – 50
<i>markewitschi</i> (Kowal, 1949) (<i>Allocreadium</i>)	D – 37
<i>martinovici</i> Ergens, 1970 (<i>Dactylogyrus</i>)	M – 94
<i>martis</i> Gmelin, 1790 (<i>Filaria</i>)	N – 400
<i>masu</i> (Fujita, 1940) (<i>Philometroides</i>)	N – 285
<i>mathevossianae</i> (Kurashvili, 1950) (<i>Diploposthe</i>)	C – 130
<i>maternus</i> Bray et Justine, 2006 (<i>Proisorhynchus</i>)	D – 596
<i>mathevossianae</i> (Schatakhstinskaya, 1953) (<i>Shiginella</i>)	D – 588
<i>matovi</i> Ergens et Kakacheva-Avramova, 1966 (<i>Gyrodactylus</i>)	M – 169
<i>mcintoshii</i> Price, 1936 (<i>Ascocotyle</i>) mtc.	D – 399
<i>mcintoshii</i> Price, 1936 (<i>Ascocotyle</i>)	D – 447
<i>mclennanae</i> (Brooks et Amato, 1992) (<i>Rhinebothroides</i>)	C – 395
<i>mediacanthus</i> Achmerow, 1952 (<i>Ancylo-discoides</i>)	M – 111
<i>medians</i> Olsson, 1876 (<i>Pleurogenoides</i>)	D – 74
<i>medinensis</i> (Linnaeus, 1758) (<i>Dracunculus</i>)	N – 103, N - 824
<i>mediovitellata</i> (Tsimbaliuk and Roytman, 1966) (<i>Allobunodera</i>)	D – 29
<i>medius</i> (Linton, 1908) (<i>Gorgorhynchus</i>)	A – 59
<i>medius</i> Kathariner, 1893 (<i>Gyrodactylus</i>)	M – 170
<i>meekii</i> Mendoza-Franco, Scholz et Vidal-Martínez, 1997 (<i>Sciadicleithrum</i>)	M – 347
<i>megacanthum</i> Ivanov et Campbell, 1997 (<i>Echinobothrium</i>)	C – 288
<i>megacephalum</i> Diesing, 1850 (<i>Zygobothrium</i>)	C – 401
<i>megalocephala</i> Price, 1932 (<i>Ascocotyle</i>) mtc.	D – 351
<i>megalocephala</i> Price, 1932 (<i>Ascocotyle</i>)	D – 446
<i>megalodiscus</i> (<i>Sciadocephalus</i>)	C – 256
<i>megalopsis</i> Rakotofitinga et Oliver, 1987 (<i>Diplectanocotyla</i>)	M - 384
<i>megalops</i> (Nitzsch et Creplin, 1829) (<i>Cloacotaenia</i>)	C – 83
<i>megalorchis</i> Nybelin, 1926 (<i>Phyllodistomum</i>)	D – 121
<i>megastomus</i> (Baer, 1944) (<i>Neoglyphe</i>)	D – 330

<i>mehrai</i> Pande, 1937 (<i>Diplodiscus</i>)	D – 300
<i>mehrai</i> Pande et Shukla, 1976 (<i>Haplorchoides</i>)	D – 267
<i>mehrai</i> Pande et Shukla, 1976 (<i>Haplorchoides</i>) mtc.	D – 276
<i>mehransis</i> Srivastava, 1933 (<i>Halipegus</i>)	D – 231
<i>mekongensis</i> Scholz, 1993 (<i>Pleurogenoides</i>)	D – 302
<i>melanocephala</i> (Rudolphi, 1819) (<i>Oncophora</i>)	N – 712
<i>melanopteri</i> Řehulková et Gelnar, 2006 (<i>Dactylogyrus</i>)	M – 226
<i>melis</i> (Schrank, 1788) (<i>Euparyphium</i>)	D – 127
<i>menschikowi</i> Gvosdev, 1950 (<i>Gyrodactylus</i>)	M – 167
<i>mentulatus</i> (Rudolphi, 1819) (<i>Plagiorchis</i>)	D – 104
<i>mergi</i> Dubois, 1932 (<i>Diplostomum</i>)	D – 562
<i>meridionalis</i> Bychowsky, 1936 (<i>Dactylogyrus</i>)	M – 98
<i>meridionalis</i> Kuperman, 1968 (<i>Triaenophorus</i>)	C – 433
<i>merlangi</i> (Kohn in Nordmann, 1832) (<i>Diclidophora</i>)	M – 400
<i>merulae</i> Baylis, 1927 (<i>Syngamus</i>)	N – 380
<i>merus</i> Zaika, 1961 (<i>Dactylogyrus</i>)	M – 88
<i>mesostoma</i> (Rudolphi, 1803) (<i>Brachylaima</i>)	D – 594
<i>metoecus</i> (Braun, 1900) (<i>Crepidostomum</i>)	D – 42
<i>mexicana</i> Moravec, Salgado-Maldonado et mayen-Peña, 1996 (<i>Aleuris</i>)	N – 652
<i>mexicana</i> Moravec, Salgado-Maldonado et Vivas-Rodríguez, 1995 (<i>Ascarophis</i>)	N – 644
<i>mexicana</i> Moravec et Fajer-Avila, 2000 (<i>Huffmanella</i>)	N – 759
<i>mexicana</i> Caspeta-Mandujano, Moravec et Salgado-Maldonado, 2000 (<i>Rhabdochona</i>)	N – 758
<i>mexicanum</i> Kritsky, Vidal-Martínez et Rodríguez-Canue, 1994 (<i>Sciadicleithrum</i>)	M - 381
<i>mexicanus</i> Caspeta-Mandujano, Moravec et Aguilar-Aguilar, 2001 (<i>Cucullanus</i>)	N – 767
<i>mexicanus</i> Caspeta-Mandujano, Moravec et Salgado-Maldonado, 1999 (<i>Dichelyne</i>)	N – 728
<i>mexicanus</i> Moravec, Salgado-Maldonado et Caspeta-Mandujano, 2000 (<i>Procamallanus</i>)	N – 722
<i>mexicanum</i> Kritsky, Vidal-Martínez et Rodríguez-Canul, 1994 (<i>Sciadicleithrum</i>)	M - 343
<i>mexicanus</i> Caspeta-Mandujano, Moravec et Salgado-Maldonado, 1999 (<i>Spinitectus</i>)	N – 747
<i>micipsae</i> (Seurat, 1917) (<i>Parapharyngodon</i>)	N – 185
<i>micracanthus</i> Nybelin, 1937 (<i>Dactylogyrus</i>)	M – 46
<i>micracanthus</i> Hukuda, 1940 (<i>Gyrodactylus</i>)	M – 197
<i>micrancristrota</i> (Wedl, 1855) (<i>Hymenolepis</i>)	C – 139
<i>microcephalum</i> (Rudolphi, 1809) (<i>Contraeaecum</i>)	N – 145
<i>microcephalus</i> (Rudolphi, 1819) (<i>Anchistrocephalus</i>)	C – 195
<i>microcephalus</i> (Dujardin, 1845) (<i>Serpinema</i>)	N – 133
<i>micropapillatum</i> (Stossich, 1890) (<i>Contraeaecum</i>)	N – 236
<i>micropapillatus</i> Törnquist, 1931 (<i>Cucullanus</i>)	N – 563
<i>micropeltis</i> Lim et Furtado, 1985 (<i>Sundanonchus</i>)	M – 279
<i>microphalos</i> (Krabbe, 1969) (<i>Anomotaenia</i>)	C – 70
<i>microphagus</i> Ingles, 1936 (<i>Megalodiscus</i>)	D – 5
<i>microrhyncha</i> (Krabbe, 1869) (<i>Anomotaenia</i>)	C – 71

<i>microscopicus</i> Woodland, 1935 (<i>Proteocephalus</i>)	C - 408
<i>microstoma</i> (Drasche, 1884) Seurat, 1918 (<i>Mehdiella</i>)	N - 335
<i>microstomum</i> Denton et Byrd, 1951 (<i>Lutztrema</i>)	D - 157
<i>midoriensis</i> Shimazu, 1990 (<i>Proteocephalus</i>)	C - 362
<i>mikhailovi</i> Ergens et Ibragimov, 1976 (<i>Gyrodactylus</i>)	M - 226
<i>militaris</i> (Rudolphi, 1803) (<i>Himasthla</i>)	D - 571
<i>minima</i> Moravec et Daniel, 1976 (<i>Rhabdochona</i>)	N - 109
<i>minima</i> (Stunkard, 1938) (<i>Stunkardiella</i>)	D - 313
<i>minima</i> (Stunkard, 1938) (<i>Stunkardiella</i>) mtc.	D - 333
<i>minimum</i> (MacCallu, 1921) (<i>Posthodiplostomum</i>) mtc.	D - 327
<i>minor</i> Wagener, 1857 (<i>Dactylogyrus</i>)	M - 47
<i>minor</i> (Steward, 1914) (<i>Spinitectus</i>)	N - 272
<i>minuta</i> (Coil, 1950) (<i>Valipora</i>)	C - 240
<i>minutum</i> (Yorke et Macfie, 1918) (<i>Trichonema</i>)	N - 547
<i>minutus</i> Kulwiec, 1927 (<i>Dactylogyrus</i>)	M - 35
<i>minutus</i> (Rudolphi, 1819) (<i>Dichelyne</i>)	N - 115
<i>minutus</i> (Goeze, 1782) (<i>Polymorphus</i>) larv.	A - 54
<i>misgurni</i> Ling, 1962 (<i>Gyrodactylus</i>)	M - 165
<i>miyatai</i> Saito, Chai, Kim, Lee et Rim, 1997 (<i>Metagonimus</i>) mtc.	D - 555
<i>mogurndae</i> (Yamaguti et Miyata, 1940) (<i>Amurotaenia</i>)	C - 371
<i>molgeta</i> Lewis, 1928 (<i>Oswaldocruzia</i>)	N - 231
<i>molini</i> Lent et Freitas, 1946 (<i>Plagiorchis</i>)	D - 105
<i>mollienisicola</i> Sogandares-Bernal et Bridgman, 1960 (<i>Ascocotyle</i>)	D - 450
<i>molnari</i> Ergens, 1978 (<i>Gyrodactylus</i>)	M - 240
<i>molnari</i> Ergens et Dulmaa, 1969 (<i>Dactylogyrus</i>)	M - 77
<i>momplei</i> Dollfus, 1932 (<i>Plagiorchis</i>)	D - 103
<i>monas</i> (Rudolphi, 1819) (<i>Mesocoelium</i>)	D - 290
<i>monenteron</i> (Wagener, 1857) (<i>Tetraonchus</i>)	M - 261
<i>monenteron</i> (Price et McIntosh, 1935) (<i>Lutztrema</i>)	D - 159
<i>mongulicus</i> Ergens et Dulmaa, 1970 (<i>Gyrodactylus</i>)	M - 203
<i>monopteri</i> Moravec et Wang, 2002 (<i>Philometra</i>)	N - 787
<i>montana</i> Yamaguti, 1943 (<i>Syphacia</i>)	N - 250
<i>montanus</i> Bychowsky, 1957 (<i>Gyrodactylus</i>)	M - 221
<i>montschadskiyi</i> Gussev, 1955 (<i>Dactylogyrus</i>)	M - 82
<i>moraveci</i> Esteban, Mas-Coma, Oltra-Ferrero et Botella, 1991 (<i>Aonchotheca</i>)	N - 565
<i>moraveci</i> Ergens, 1979 (<i>Gyrodactylus</i>)	M - 248
<i>moraveci</i> Richardson et Adamson, 1988 (<i>Megalobatrachonema</i>)	N - 244
<i>moraveci</i> Baruš, Kajerová et Koubková, 2005 (<i>Pterothominx</i>)	N - 840
<i>moravicus</i> Sitko, 1993 (<i>Plagiorchis</i>)	D - 585
<i>mrazeki</i> Ergens et Dulmaa, 1969 (<i>Dactylogyrus</i>)	M - 78
<i>mucronata</i> (Molin, 1858) (<i>Pearsonema</i>)	N - 329
<i>mugilis</i> Liu et Yang, 2002 (<i>Haploporus</i>)	D - 456
<i>muehlingi</i> (Jagerskiold, 1899) (<i>Apophallus</i>)	D - 437
<i>multiglandularis</i> Semenow, 1927 (<i>Plagiorchis</i>)	D - 474
<i>multipapillatum</i> (Drasche, 1882) (<i>Contracaecum</i>)	N - 425
<i>multistriatus</i> (Rudolphi, 1810) (<i>Colymbilepsis</i>)	C - 84
<i>muris</i> (Schränk, 1788) (<i>Trichocephalus</i>)	N - 440
<i>muris</i> (Gmelin, 1790) (<i>Mastophorus</i>)	N - 92
<i>murissylvatici</i> (Diesing, 1851) (<i>Aonchotheca</i>)	N - 316
<i>musculara</i> (Hart, 1936) (<i>Grillotia</i>)	C - 11

<i>musculosa</i> (Fuhrmann, 1896) (<i>Choanotaenia</i>)	C – 144
<i>musteli</i> (Yamaguti, 1952) (<i>Erudituncus</i>)	C – 334
<i>musteli</i> (Beneden, 1850) (<i>Orygmatobothrium</i>)	C – 277
<i>mutabile</i> (Zeder, 1800) (<i>Cyclocoelum</i>)	D – 198
<i>mutabilis</i> (Rudolphi, 1819) (<i>Anomotaenia</i>)	C – 72
<i>mutabilis</i> (Rudolphi, 1819) (<i>Polymorphus</i>)	A – 22
<i>mutabilis</i> (Rudolphi, 1819) (<i>Hexaglandula</i>)	N – 441
<i>mutabilitas</i> Bychowsky, 1957 (<i>Gyrodactylus</i>)	M – 233
<i>mutabilis</i> Linton, 1927 (<i>Valipora</i>)	C – 302
<i>mühlingi</i> (Jägerskiöld, 1899) (<i>Apophallus</i>)	D – 140
<i>myiophoneii</i> Baruš, Kullmann et Tenora, 1972 (<i>Thelaziella</i>)	N – 355
<i>myliobatidis</i> Shipley et Hornell, 1906 (<i>Rhoptrobothrium</i>)	C – 430
<i>mysti</i> Karve, 1952 (<i>Procamallanus</i>)	N – 196
<i>nagibinae</i> (Gläser, 1965) (<i>Paradiplozoon</i>)	M – 302
<i>najal</i> De, 1997 (<i>Paracapillaria</i>)	N – 688
<i>nana</i> Ranson, 1920 (<i>Ascocotyle</i>)	D – 340
<i>nana</i> Ranson, 1920 (<i>Ascocotyle</i>) mtc.	D – 341
<i>nanaflexicollis</i> Dranen, Rubec et Underwood, 1977 (<i>Bathycreadium</i>)	D – 494
<i>nanoides</i> Gussev, 1966 (<i>Dactylogyrus</i>)	M – 21
<i>nanus</i> (Rudolphi, 1802) (<i>Plagiorchis</i>)	D – 475
<i>nanus</i> Dogiel et Bychowsky, 1934 (<i>Dactylogyrus</i>)	M – 48
<i>narzikulovi</i> Ergens et Dzhililov, 1979 (<i>Gyrodactylus</i>)	M – 246
<i>nasalis</i> Ergens, 1988 (<i>Paraquadriacanthus</i>)	M – 323
<i>nasalis</i> Strelkow et Ha Ky, 1964 (<i>Dactylogyrus</i>)	M – 27
<i>nasaatum</i> (Loos, 1900) Ihle, 1922 (<i>Cylicocycclus</i>)	N – 245
<i>nasuta</i> (Rudolphi, 1819) (<i>Dispharynx</i>)	N – 221
<i>nattereri</i> Parona, 1901 (<i>Ophiotaenia</i>)	C – 226
<i>naviculoides</i> Ergens, 1956 (<i>Dactylogyrus</i>)	M – 62
<i>nayaritensis</i> Caspeta-Mandujano, Moravec, Salgado-Maldonado, 2000 (<i>Fellicola</i>)	N – 760
<i>neglecta</i> (Diesing, 1851) (<i>Icosiella</i>)	N – 89
<i>neglectus</i> La Rue, 1911 (<i>Proteocephalus</i>)	C – 31
<i>nemachili</i> Dubinina, 1957 (<i>Digramma</i>)	C – 424
<i>nemachili</i> Bychowsky, 1936 (<i>Gyrodactylus</i>)	M – 208
<i>nemachilus</i> (Kaw, 1950) (<i>Allocreadium</i>)	D – 38
<i>nemipteri</i> Moravec et Shahaarom, 1989 (<i>Paraphilometroides</i>)	N – 372
<i>nemipteri</i> Moravec et Justine, 2005 (<i>Raphidascaris</i>)	N – 836
<i>neocaballeri</i> (<i>Procamallanus</i> – <i>Spirocamallanus</i>)	N – 651
<i>neocomense</i> Fuhrmann, 1927 (<i>Petasiger</i>)	D – 584
<i>nicaraguense</i> Vidal-Martínez, Scholz et Aguirve-Macedo, 2001 (<i>Sciadicleithrum</i>)	M – 363
<i>nicholsii</i> Bray Gibson, 1980 (<i>Prudhoeus</i>)	D – 539
<i>nigrescens</i> Mendoza-Franco, Violante-González et Vidal-Martínez, 2006 (<i>Cornutohaptor</i>)	M – 418
<i>nigrovenosus</i> (Bellingham, 1844) (<i>Leptophallus</i>)	D – 75
<i>nimia</i> Linton, 1910 (<i>Helicometrina</i>)	D – 369
<i>nipponensis</i> Ogawa et Egusa, 1978 (<i>Gyrodactylus</i>)	M – 213
<i>nipponensis</i> Yamaguti, 1941 (<i>Raphidascaroides</i>)	N – 734
<i>nipponica</i> Yamaguti, 1935 (<i>Rhabdias</i>)	N – 162
<i>nobilis</i> (Long et Yu, 1958) (<i>Dactylogyrus</i>)	M – 1

<i>nodulosa</i> Mackiewicz et McCrae, 1962 (<i>Hunterella</i>)	C – 321
<i>nodulosus</i> (Pallas, 1781) (<i>Triaenophorus</i>) plerocerkoid	C - 40 C – 28
<i>nokomis</i> Macy, 1937 (<i>Prosthodendrium</i>)	D – 160
<i>nordiana</i> Bashkirova, 1946 (<i>Echinostoma</i>)	D – 601
<i>nordmanni</i> Ergens et Dulmaa, 1970 (<i>Gyrodactylus</i>)	M – 202
<i>norvegia</i> Moravec et Karlsbakk, 2000 (<i>Capillostrongyloides</i>)	N – 757
<i>notacanthi</i> Bray, 2004 (<i>Steringovermes</i>)	D – 543
<i>notopteri</i> (Jain, 1955 Lim et Furtado, 1986 (<i>Silurodiscoides</i>))	M – 321
<i>nouveli</i> Seurat, 1914 (<i>Tropisurus</i>)	N – 297
<i>novaeguineae</i> Moravec et Sey, 198 (<i>Cosmocerca</i>)	N – 387
<i>novaezelandiae</i> Moravec et Taraschewski (<i>Anguillicola</i>)	N – 263
<i>numidica</i> (Seurat, 1917) Sprent, 1985 (<i>Seuratascaris</i>)	N – 392
<i>numidica</i> Linstow, 1899 (<i>Oxynema</i>)	N – 305
<i>nunezae</i> Scholz, Vargas-Vázquez, Vidal-Martínez et Aguirre-Macedo, 1996 (<i>Ascocotyle</i>)	D – 348
<i>nunezae</i> Scholz, Vargas-Vázquez, Vidal-Martínez et Aguirre-Macedo, 1996 mtc. (<i>Ascocotyle</i> A.)	D – 349
<i>nursei</i> Ergens, 1973 (<i>Characidotrema</i>)	M – 282
<i>nyrocae</i> (Yamaguti, 1935) (<i>Diorchis</i>)	C – 121
<i>obesa</i> (Diesing, 1850) (<i>Prosthenhystera</i>)	D – 383
<i>obesum</i> Loos, 1902 (<i>Saccocelium</i>)	D – 14
<i>oblonga</i> Cobbold, 1858 (<i>Campula</i>)	D – 409
<i>oblongus</i> (Wilson, 1908) (<i>Perissopus</i>)	Cr – 5
<i>obscurum</i> (Leidy, 1887) (<i>Cyclocoelum</i>)	D – 515
<i>obsignata</i> Madsen, 1945 (<i>Capillaria</i>)	N – 505
<i>obsignata</i> (Madsen, 1945) (<i>Baruscapillaria</i>)	N – 153
<i>obturans</i> (Prenant, 1886) (<i>Philometra</i>)	N – 62
<i>obtusa</i> Rudolphi, 1802 (<i>Diplotriaena</i>)	N – 309
<i>obtusum</i> (Looss, 1896) (<i>Prosthodendrium</i>)	D – 102
<i>obtusus</i> Dujardin, 1845 (<i>Proleptus</i>)	N – 762
<i>obvelata</i> Rudolphi, 1802 (<i>Syphacia</i>)	N – 211
<i>obvelatus</i> Creplin, 1825 (<i>Cosmocephalus</i>)	N – 299
<i>oceaniensis</i> Moravec, Sasal, Würtz et Taraschewski, 2005 (<i>Cucullanus</i>)	N – 844
<i>octacantha</i> (Krabbe, 1869) (<i>Dicranotaenia</i>)	C – 88
<i>octacantha</i> (Krabbe, 1869) (<i>Sobolevicanthus</i>)	C – 185
<i>octohamatus</i> Glennon, Chisholm et Whittington, 2005 (<i>Branchotenthes</i>)	M – 412
<i>ocularis</i> Moravec et Ogawa, 2002 (<i>Philometra</i>)	N – 782
<i>oculeus</i> Kossack, 1911 (<i>Hyptiasmus</i>)	D – 519
<i>oesophagea</i> (Polonio, 1859) (<i>Dracunculus</i>)	N – 129
<i>oesophagicola</i> Soltys, 1952 (<i>Eucoleus</i>)	N – 274
<i>okavangoensis</i> Moravec et Van As, 2004 (<i>Synodontisia</i>)	N – 821
<i>oligoon</i> (Linstow, 1887) (<i>Psilotrema</i>)	D – 613
<i>olrikii</i> Krabbe, 1874 (<i>Diplocotyle</i>)	C – 333
<i>olssoni</i> (Railliet, 1900) (<i>Brachylecithum</i>)	D – 433
<i>oncorhynchi</i> (Fujita, 1921) (<i>Rhabdochona</i>)	N – 44
<i>oncorhynchi</i> Ogawa et Egusa, 1978 (<i>Tetraonchus</i>)	M – 278
<i>oncorhynchi</i> Kuitunen-Ekbaum, 1933 (<i>Philonema</i>)	N – 63
<i>ondatrae</i> (Price, 1931) (<i>Ribeiroia</i>)	D – 292

<i>opaca</i> (Wisniewski, 1934) (<i>Cyathocotyle</i>)	D – 560
<i>opeatura</i> Leidy, 1891 (<i>Atractis</i>)	N – 711
<i>ophiocephali</i> (Thapar, 1931) (<i>Pallisentis</i>)	A – 32
<i>ophicephali</i> (Pearse, 1933) (<i>Neocamallanus</i>)	N – 194
<i>ophicephali</i> Ha Ky, 1971 (<i>Spinitectus</i>)	N – 201
<i>ophiocephali</i> Gussev, 1955 (<i>Gyrodactylus</i>)	M – 201
<i>ophiosterni</i> Moravec, Salgado-Maldonado et Caspeta-Mandujano, 1992 (<i>Pseudocapillaria</i>)	N – 748
<i>ophisauri</i> (Kreis, 1940) (<i>Entomelas</i>)	N – 371
<i>ophisterni</i> Scholz, de Chambrier et Salgado-Maldonado, 2001 (<i>Monticellia</i>)	C – 335
<i>ophisterni</i> Moravec, Salgado-Maldonado et Aguilar-Aguilar, 2002 (<i>Philometra</i>)	N – 781
<i>ophiusae</i> Roca et Hornero, 1992 (<i>Strongyloides</i>)	N – 593
<i>opisthovitellinum</i> (Soltys, 1954) (<i>Rubensstrema</i>)	D – 329
<i>orbiculare</i> (Stunkard, 1916) (<i>Neopolystoma</i>)	M – 3
<i>orensensis</i> Cremonte et Martoreli, 1998 (<i>Maritrema</i>)	D – 405
<i>oreoleucisci</i> Ergens et Dulmaa, 1970 (<i>Dactylogyrus</i>)	M – 76
<i>oreoleucisci f. typica</i> Ergens et Dulmaa, 1970 (<i>Gyrodactylus</i>)	M – 204
<i>oreoleucisci</i> Moravec et Ergens, 1970 (<i>Philometra</i>)	N – 8
<i>orientalis</i> Kulakovskaja, 1962 (<i>Braviscolex</i>)	C – 219
<i>orientalis</i> Heller, 1868 (<i>Nessipus</i>)	Cr – 6
<i>orientalis</i> Kuperman, 1968 (<i>Triaenophorus</i>)	C – 433
<i>orientalis</i> Kuperman, 1968 larv. (<i>Triaenophorus</i>)	C – 435
<i>ornata japonica</i> Yamaguti (<i>Cosmocerca</i>)	N – 456
<i>ornata</i> (Dujardin, 1845) (<i>Cosmocerca</i>)	N – 86
<i>ornatohamus</i> Lim et Furtado, 1986 (<i>Notopterodiscooides</i>)	M – 318
<i>orphoidis</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 296
<i>osculatum</i> (Rudolphi, 1802) (<i>Contracaecum</i>)	N – 234
<i>osculatus</i> (Goeze, 1782) (<i>Proteocephalus</i>)	C – 49
<i>osheroffi</i> Meggitt, 1934 (<i>Oochoristica</i>)	C – 249
<i>oschmarini</i> Trofimenko (<i>Comephoronema</i>)	N – 721
<i>osoblahensis</i> Ergens, 1963 (<i>Gyrodactylus</i>)	M – 185
<i>osorioi</i> Choudhury et Pérez-Ponce de León, 2001 (<i>Spinitectus</i>)	N – 802
<i>ossicola</i> Justine, 2004 (<i>Huffmanella</i>)	N – 818
<i>osteochili</i> (Lim et Furtado, 1984) (<i>Dactylogyrus</i>)	M – 288
<i>otsurui</i> Saito et Shimizu, 1968 (<i>Metagonimus</i>) mtc.	D – 554
<i>ovale</i> (Linstow, 1907) (<i>Contracaecum</i>)	N – 515
<i>ovale</i> Martin, 1950 (<i>Phocitrema</i>)	D – 395
<i>ovata</i> (Zeder, 1803) (<i>Philometra</i>)	N – 84
<i>ovatus</i> Katsurada, 1914 (<i>Paracoenogonimus</i>)	D – 579
<i>ovatus</i> (Rudolphi, 1803) (<i>Prosthogonimus</i>)	D – 136
<i>overstreeti</i> Deardorff et Ko, 1983 (<i>Echinocephalus</i>)	N – 848
<i>overstreeti</i> Moravec et de Buron, 2006 (<i>Philometra</i>)	N – 857
<i>ovifilamenta</i> Moravec et Thatcher, 2001 (<i>Ichthyouris</i>)	N – 778
<i>oviflagellis</i> Fourment, 1883 (<i>Spinitectus</i>)	N – 851
<i>ovis</i> Orlov, Ershov et Badanin, 1934 (<i>Skrjabinotrema</i>)	D – 131
<i>ovis</i> (Cobbold, 1869) (<i>Taenia</i>)	C – 248
<i>ovofurcata</i> Czaplinski, 1972 (<i>Diorchis</i>)	C – 123
<i>ovolaciniata</i> (Linstow, 1877) (<i>Anomotaenia</i>)	C – 54
<i>ovopunctata</i> (Linstow, 1873) (<i>Baruscapillaria</i>)	N – 225
<i>oxycephala</i> (<i>Sebekia</i>)	L – 2

<i>oxycephalus</i> Ward et Magath, 1917 (<i>Camallanus</i>)	N – 10
<i>oxyurus</i> (Creplin, 1825) (<i>Psilochasmus</i>)	D – 504
<i>ozakii</i> Nagaty, 1937 (<i>Bucephalopsis</i>)	D – 210
<i>pacifica</i> Moravec, Salgado-Maldonado et Caspeta-Mandujano, 1999 (<i>Paracapillaria teixeirafreitasi</i>)	N – 726
<i>pacificus</i> Moravec, Justine, Würtz, Taraschewski et Sasal, 2005	N – 843
<i>pahangensis</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 289
<i>pachystoma</i> (Eysenhardt, 1829) (<i>Haplospalchnus</i>)	D – 8
<i>pallidum</i> Stafford, 1904 (<i>Homalometra</i>)	D – 307
<i>panamense</i> Caballero, Bravo-Hollis et Grocott, 1953 (<i>Pseudacanthostomum</i>)	D – 384
<i>panamense</i> Caballero, Bravo-Hollis et Grocott, 1953 (<i>Pseudoacanthostomum</i>)	D – 420
<i>pancreaticum</i> (Janson, 1889) (<i>Eurytrema</i>)	D – 248
<i>pancreaticus</i> Baer, 1960 (<i>Euamphimerus</i>)	D – 531
<i>paniceum</i> Brinkmann, 1942 (<i>Pachytrema</i>)	D – 187
<i>panonicus</i> Molnár, 1968 (<i>Gyrodactylus</i>)	M – 127
<i>papernai</i> Moravec et Taraschewski, 1988 (<i>Anguillicola</i>)	N – 279
<i>papilligerum</i> (Rees, 1968) (<i>Allocreadium</i>)	D – 39
<i>papilocirratum</i> Groschaft et Sitko, 1970 (<i>Leucochloridium</i>)	D – 193
<i>papillosus</i> Wedl, 1856 (<i>Strongyloides</i>)	N – 451
<i>papuensis</i> Moravec et Sey, 198 (<i>Desmognathinema</i>)	N – 286
<i>parablennii</i> Moravec, Orecchia et Paggi, 1988 (<i>Pseudocapillaria</i>)	N – 283
<i>paracinctum</i> Bychowskaja-Pawlovskaja, 1954 (<i>Echinoparyphium</i>)	D – 149
<i>paracompressa</i> (Czaplinski, 1956) (<i>Microsomacanthus</i>)	C – 159
<i>paradoxum</i> (Nordmann, 1832) (<i>Diplozoon</i>)	M – 298
<i>paradoxum</i> Carus, 1833 (<i>Leucochloridium</i>)	D – 573
<i>paradoxus</i> Creplin, 1839 (<i>Ancyrocephalus</i>)	M – 103
<i>parafimbriatum</i> Befus et Freeman, 1973 (<i>Corallobothrium</i>)	C – 234
<i>paralichthydis</i> Moravec et de Buron, 2006 (<i>Philometroides</i>)	N – 858
<i>paramicrosoma</i> (Gasowska, 1932) (<i>Microsomacanthus</i>)	C – 156
<i>paraminimus</i> (Ergens, 1966) (<i>Gyrodactylus</i>)	M – 336
<i>paranaensis</i> (Pavanelli et Rego, 1989) (<i>Chambriella</i>)	C – 401
<i>paranemachili</i> Ergens et Bychowsky, 1967 (<i>Gyrodactylus</i>)	M – 199
<i>parasiluri</i> Yamaguti, 1934 (<i>Gangesia</i>)	C – 125
<i>parasiluri</i> Yamaguti, 1934 (<i>Gangesia</i>) larv.	C – 352
<i>parasiluri</i> Yin et Wu, 1984 (<i>Heterosentis</i>)	A – 74
<i>parasiluri</i> (Zmeev, 1936) (<i>Paraproteocephalus</i>)	C – 367
<i>paronai</i> Moravec et Garibaldi, 2000 (<i>Huffmanella</i>)	N – 763
<i>parophrysi</i> (Moravec et al., 1981) (<i>Paracapillaria</i>)	N – 3
<i>partipentazonae</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 297
<i>parupenei</i> Moravec, Orecchia et Paggi, 1988 (<i>Ascarophis</i>)	N – 251
<i>parva</i> (Zmeev, 1936) (<i>Khawia</i>)	C – 339
<i>parviuncinatum</i> Young, 1954 (<i>Acanthobothrium</i>)	C – 319
<i>parviventosom</i> Dubois, 1932 (<i>Diplostomum</i>)	D – 563
<i>parvogenitalis</i> (Mathevossian, 1946) (<i>Diorchis</i>)	C – 124
<i>parvus</i> Bychowsky, 1936 (<i>Gyrodactylus</i>)	M – 231
<i>parvus</i> Wegener, 1910 (<i>Dactylogyrus</i>)	M – 53
<i>passerina</i> (Fuhrmann, 1907) (<i>Choanotaenia</i>)	C – 146
<i>passerina</i> (Fuhrmann, 1907) (<i>Anomotaenia</i>)	C – 55
<i>passerina</i> Fuhrmann, 1908 (<i>Biuterina</i>)	C – 95

<i>passeris</i> (Gmelin, 1790) (<i>Passerilepis</i>)	C – 193
<i>passeris</i> (Gmelin, 1790) (<i>Dicranotaenia</i>)	C – 89
<i>patagonense</i> Moravec et Urawa, 1996 (<i>Hysterothylacium</i>)	N – 675
<i>patagonicum</i> Shimazu, 1999 (<i>Allocreadium</i>)	D – 424
<i>patagonicus</i> (Szidat, 1956) (<i>Deropegus</i>)	D – 425
<i>pateratum</i> (Yorke et Macfie, 1919) (<i>Cylicodontophorus</i>)	N – 545
<i>patzcuarensis</i> Osorio-Sarabia, Pérez-Ponce de León et Salgado-Maldonado, 1986 (<i>Capillaria</i>)	N – 718
<i>paucispinosum</i> Labriola et Suriano, 2000 (<i>Wardium</i>)	C – 311
<i>paucispinus</i> (Sandground, 1928) (<i>Tropisurus</i>)	N – 220
<i>pavlovskii</i> (Bychowsky et Nagibina, 1959) (<i>Diplozoon</i>)	M – 378
<i>pavlovskii</i> Ergens et Bychowsky, 1967 (<i>Gyrodactylus</i>)	M – 159
<i>pavlovskii</i> Dubinina, 1959 (<i>Ligula</i>)	C – 372
<i>pavlovskii</i> (Bychowsky et Nagibina, 1959) (<i>Paradiplozoon</i>)	M – 304
<i>pavlovskii</i> Bychowsky et Gussev, 1955 (<i>Pseudacolpenteron</i>)	M – 283
<i>pawlovskii</i> (Zmeev, 1936) (<i>Phyllodistomum</i>)	D – 463
<i>pearsei</i> Scholz, Vargas-Vázquez, Moravec, 1996 (<i>Bothriocephalus</i>)	C – 244
<i>pectinata</i> (Goeze, 1782) (<i>Mosgovoyia</i>)	C – 385
<i>pectinatus</i> Kuchta et Scholz, 2005 (<i>Australicola</i>)	C – 384
<i>peekayi</i> Cribb et Pichelin, 1997 (<i>Notopronocephalus</i>)	D – 377
“ <i>pekinensis</i> “ Groschaft (?) (<i>Brachylecithum</i>)	D – 174
<i>pelargopsis</i> Nandi et Mujumdar, 1988 (<i>Schistogendra</i>)	N – 151
<i>pentabrachiatus</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 295
<i>pentabrachicleithrium</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 287
<i>pentasoma</i> (Klaptocz, 1906) (<i>Proteocephalus</i>)	C – 396
<i>peraccuratus</i> Pinto, Fabio, Noronha et Rolas, 1976 (<i>Procamallanus</i>)	N – 577
<i>perakensis</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 296
<i>percae</i> Ergens, 1966 (<i>Ancyrocephalus</i>)	M – 104
<i>percae</i> (Müller, 1780) (<i>Proteocephalus</i>)	C – 29
<i>perccotti</i> Ergens et Yuhhimenko, 1973 (<i>Gyrodactylus</i>)	M – 139
<i>percichthydis</i> Moravec, Urawa et Coria, 1997 (<i>Philonema</i>)	N – 690
<i>perfectum</i> (Beneden, 1853) (<i>Monorygma</i>)	C – 275
<i>perforans</i> (Kotlan et Orosz, 1931) (<i>Eucoleus</i>)	N – 662
<i>perforator</i> (Linstow, 1908) (<i>Pomphorhynchus</i>)	A – 17
<i>perlatum</i> Clurea, 1911 (<i>Conodiplostomum</i>)	D – 597
<i>petasaurum</i> (Deslongschaps, 1824) (<i>Notocotyloides</i>)	D – 578
<i>petiolatus</i> (Railliet, 1900) (<i>Zonorchis</i>)	D – 509
<i>petkovici</i> Ergens, 1970 (<i>Dactylogyrus</i>)	M – 91
<i>petrowi</i> Belous, 1965 (<i>Spinitectus</i>)	N – 812
<i>petruschewskii</i> Sokolovskaya, 1962 (<i>Dendronucleata</i>)	A – 30
<i>petruschewskii</i> (Schulman, 1948) (<i>Schulmanella</i>)	N – 53
<i>petterae</i> Moravec et Sey, 1988 (<i>Procamallanus</i>)	N – 50
<i>petterae</i> Moravec, Kohn, Fernandes, 1991 (<i>Parasynodontisia</i>)	N – 550
<i>pezowii</i> Ergens, 1980 (<i>Gyrodactylus</i>)	M – 205
<i>phalacrocoracis</i> (Yamaguti, 1939) (<i>Andracantha</i>)	A – 53
<i>phalacrocoracis</i> (Yamaguti, 1939) (<i>Petasisiger</i>)	D – 297
<i>philactes</i> (Schiller, 1951) (<i>Parabisaccanthes</i>)	C – 104
<i>philippinensis</i> (Chitwood, Velasquez et Salazar, 1966) Moravec, 2000 (<i>Paracapillaria</i>)	N – 432, N – 756

<i>phoxini</i> Malmberg, 1957 (<i>Gyrodactylus</i>)	M – 182
<i>phoxini</i> Moravec, 1968 (<i>Rhabdochona</i>)	N – 45
<i>phoxini</i> Malewiczka, 1949 (<i>Dactylogyrus</i>)	M – 73
<i>phrynomantisi</i> Moravec, 1990 (<i>Cosmocercella</i>)	N – 445
<i>phrynosoma</i> (Ortlepp, 1922) (<i>Skrjabinoptera</i>)	N – 566
<i>phycidis</i> (Parona et Perugia, 1889) (<i>Diclidophora</i>)	M – 414
<i>physeteris</i> Baylis, 1923 (<i>Anisakis</i>)	N – 705
<i>physiculi</i> Moravec et Nagasawa, 1999 (<i>Hysterothylacium</i>)	N – 738
<i>piayae</i> (Sandgroudnd, 1929) (<i>Cyrnea</i>)	N – 424
<i>picorum</i> (Rudolphi, 1819) (<i>Capillaria</i>)	N – 512
<i>pikaeuris</i> (Akhtar, 1953) (<i>Eugenuris</i>)	N – 353
<i>pilaetus</i> (Rudolphi, 1802) (<i>Cotylurus</i>)	D – 112
<i>pilherodiae</i> Mahon, 1956 (<i>Dendrouterina</i>)	C – 241
<i>pimellodelae</i> Moravec, Kohn, Fernandes, 1991 (<i>Brasilnema</i>)	N – 549
<i>pimelodellae</i> Moravec, Kohn et Fernandes, 1993 (<i>Cucullanus</i>)	N – 583
<i>pimelodus</i> Pinto, Fabio, Noronha et Rolas, 1974 (<i>Procamallanus</i>)	N – 590
<i>pindoramensis</i>	
<i>pindoramensis</i> (Travassos, 1929) (<i>Pygidiopsis</i>) mtc.	D – 444
<i>pinguicollum</i> (Sleggs, 1927) (<i>Pinguicollum</i>)	C – 18
<i>pinguis</i> (Mehlis, 1831) (<i>Renicola</i>)	D – 587
<i>pinnai</i> Travassos, Artigas et Pereica, 1928 (<i>Cucullanus</i>)	N – 588
<i>pinnai pterodorasi</i> Moravec, Kohn et Fernandes, 1996 (<i>Cucullanus</i>)	N – 684
<i>pintoi</i> (Kohn et Fernandes, 1988) (<i>Procamallanus</i>)	N – 681
<i>piracatinga</i> (Woodland, 1935) (<i>Rudolphiella</i>)	C - 409
<i>piramutab</i> (Woodland, 1933) (<i>Proteocephalus</i>)	C – 415
<i>piriformis</i> Carfora, de Chambrier et Vaucher, 2003 (<i>Amphoteromorphus</i>)	C - 413
<i>piscatorum</i> de Chambrier and Vaucher, 1999 (<i>Mariauxiella</i>)	C - 404
<i>pisiformis</i> (Bloch, 1780) (<i>Taenia</i>)	C – 114
<i>plagiorchis</i> Ozaki, 1926 (<i>Coitocaecum</i>)	D – 454
<i>planoratus</i> Kulkarni, 1935 (<i>Procamallanus</i>)	N – 826
<i>plataleae</i> Dubinin, 1940 (<i>Hysteromorpha</i>)	D – 602
<i>platamauxili</i> Lim et Furtado, 1986 (<i>Silurodiscooides</i>)	M – 320
<i>platycephalus</i> (Creplin, 1825) (<i>Cotylurus</i>) mtc.	D – 237
<i>platycephalus</i> (Creplin, 1825) (<i>Ichthyocotylurus</i>)	D – 604
<i>platyceri</i> Hartwich et Tscherner, 1979 (<i>Ascaridia</i>)	N – 838
<i>platyrhinorum</i> Baruš, 1961 (<i>Capillaria</i>)	N – 320
<i>plecoglossi</i> Ogawa et Egusa, 1978 (<i>Gyrodactylus</i>)	M – 189
<i>plecoglossi</i> Yamaguti, 1934 (<i>Proteocephalus</i>)	C – 305
<i>plegadisi</i> Digiani, 1999 (<i>Syncuaria</i>)	N – 749
<i>pleuronectis</i> Müller, 1788 (<i>Scolex</i>)	C – 266
<i>plica</i> (Rudolphi, 1819) (<i>Pearsonema</i>)	N – 480
<i>plotosi</i> Moravec et Nagasawa, 1989 (<i>Philometra</i>)	N – 359
<i>podicipina</i> (Szymanski, 1905) (<i>Confluaria</i>)	C – 85
<i>podicipina</i> Kazicka et Niewiadomska, 1960 (<i>Tylodelphys</i>)	D – 111
<i>podicipinus</i> Baker et Vancher, 1984 (<i>Cosmocerca</i>)	N – 731
<i>podicipitis</i> (Yamaguti, 1941) (<i>Capillaria</i>)	N – 501
<i>poeyei</i> Baruš et Valle, 1967 (<i>Tricholeiperia</i>)	N – 481
<i>pollanicola</i> Gresson, 1952 (<i>Proteocephalus</i>)	C – 290
<i>polonica</i> (Kowalewski, 1895) (<i>Bilharziella</i>)	D – 528
<i>polymorphus</i> Gussev, 1955 (<i>Ancyrocephalus</i>)	M – 101

<i>polymorphus</i> Baer, 1827 (<i>Bucephalus</i>)	D – 1
<i>polypteri</i> Vassiliadès et Chevalier, 1973 (<i>Gendria</i>)	N – 829
<i>popowi</i> Markov et Bogdanov, 1963 (<i>Thelandros</i>)	N – 363
<i>porosa</i> (Rudolphi, 1810) (<i>Choanotaenia</i>)	C – 264
<i>porosa</i> (Rudolphi, 1810) (<i>Paricterotaenia</i>)	C – 171
<i>postgonoporus</i> Neiland, 1951 (<i>Macyella</i>)	D – 175
<i>praecox</i> (Krabbe, 1882) (<i>Anomotaenia</i>)	C – 56
<i>preaputiale</i> Linstow, 1889 (<i>Physaloptera</i>)	N – 339
<i>pricei</i> Mueller, 1936 (<i>Cleidodiscus</i>)	M – 116
<i>priedei</i> Bray et Merrett, 1998 (<i>Prodistomum</i>)	D – 547
<i>prima</i> Shikhobalova, 1930 (<i>Skrjabinocerca</i>)	N – 298
<i>pristis</i> (Deslongchamps, 1824) (<i>Stephanostomum</i>)	D – 497
<i>pritchardae</i> (Campbell, 1975) (<i>Steringophorus</i>)	D – 542
<i>probolurus</i> (Railliet, 1896) (<i>Trichostrongylus</i>)	N – 532
<i>procyonis</i> (Pence, 1975) (<i>Eucoleus</i>)	N – 256
<i>proglottina</i> (Davaine, 1860) (<i>Davainea</i>)	C – 358
<i>prochilodonti</i> Domingues et Boeger, 2001 (<i>Protorhinoxenus</i>)	M – 366
<i>prokopici</i> Moravec, Baruš et Ryšavý, 1987 (<i>Strongyloides</i>)	N – 173
<i>propinquus</i> (Dujardin, 1845) Meyer, 1932 (<i>Acanthocephaloides</i>)	A – 49
<i>propinquus</i> Bychowsky, 1931 (<i>Dactylogyrus</i>)	M – 57
<i>prostae</i> Ergens, 1963 (<i>Gyrodactylus</i>)	M – 179
<i>prostae</i> Molnar, 1964 (<i>Dactylogyrus</i>)	M – 16
<i>proterorhini</i> Ergens, 1967 (<i>Gyrodactylus</i>)	M – 173
<i>proximus</i> Barker, 1915 (<i>Plagiorchis</i>)	D – 507
<i>prussica</i> Mühling, 1896 (<i>Cyathocotyle</i>)	D – 153
<i>przewalskii</i> (Skrjabin, 1914) (<i>Dropanidotaenia</i>)	C – 102
<i>pseudaspiei</i> Moravec et Ergens, 1970 (<i>Philometroides</i>)	N – 9
<i>pseudoechinata</i> (Olsson, 1876) (<i>Stephanoprora</i>)	D – 135
<i>pseudonemachili</i> Ergens et Bychowsky, 1967 (<i>Gyrodactylus</i>)	M – 209
<i>pseudoplatystomae</i> Moravec, Kohn et Fernandes, 1992 (<i>Cucullanus</i>)	N – 576
<i>pseudospathaceum</i> Niewiadomska, 1984 (<i>Diplostomum</i>)	D – 287
<i>pseudostoma</i> (Wilemoes-Suhm, 1870) (<i>Crocodilicola</i>)	D – 382
<i>pterothylli</i> Heinze, 1933 (<i>Capillaria</i>)	N – 49
<i>ptychocheilus</i> (Faust, 1917) (<i>Ornithodiplostomum</i>)	D – 33
<i>pudendotectus</i> (Wostokow, 1905) (<i>Metastrongylus</i>)	N – 475
<i>pudica</i> Seurat, 1914 (<i>Thubunaea</i>)	N – 186
<i>pulchella</i> (Rudolphi, 1819) (<i>Helicometra</i>)	D – 263
<i>pulverulenta</i> (Braun, 1901) (<i>Dendritobilharzia</i>)	D – 561
<i>pumilio</i> (Looss, 1896) (<i>Haplorchis</i>)	D – 256
<i>pumilio</i> (Looss, 1896) (<i>Haplorchis</i>) mtc.	D – 279
<i>punctatus</i> Moravec et Scholz, 1991 (<i>Procamallanus</i>)	N – 555
<i>pungens</i> (Schneider, 1866) (<i>Diplotriaena</i>)	N – 342
<i>pungens</i> (Linstow, 1884) (<i>Petasiger</i>)	D – 610
<i>pungiti</i> Shigin, 1965 (<i>Diplostomum</i>)	D – 599
<i>pungitii</i> Malmberg, 1964 (<i>Gyrodactylus</i>)	M – 239
<i>pungitii</i> Dubinina, 1959 (<i>Schistocephalus</i>)	C – 425
<i>punti</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 286
<i>pusilla</i> (Goeze, 1782) (<i>Catenotaenia</i>)	C – 391
<i>pusillum</i> Dubois, 1928 (<i>Diplostomum</i>)	D – 600
<i>putorii</i> (Rudolphi, 1819), López-Neyra, 1947	N – 315

<i>puylaerti</i> Moravec, 1983 (<i>Rhabdochona</i>)	N – 142
<i>pythonis</i> Blainville, 1824 (<i>Bothridium</i>)	C – 215
<i>quadratum</i> (Linstow, 1892) (<i>Diphyllobothrium</i>)	C – 328
<i>quadricornis</i> (Wedl, 1862) (<i>Hexametra</i>)	N – 128
<i>quadriradiatus</i> (Stevenson, 1904) (<i>Ornithostrongylus</i>)	N – 357
<i>queenslandensis</i> Jones et Beveridge, 1999 (<i>Nybelinia</i>)	C – 301
<i>quentini</i> Baruš, 1972 (<i>Helminthoxys</i>)	N – 377
<i>quindecima</i> Chisholm et Whittington, 1999 (<i>Empruthotrema</i>)	M – 352
<i>quineserialis</i> Barker et Laughlin, 1911 (<i>Quineserialis</i>)	D – 283
<i>quiscali</i> Baruš, 1968 (<i>Strongyloides</i>)	N – 454
<i>raabei</i> (Bezubik, 1958) (<i>Cotylurus</i>)	D – 520
<i>rabuka</i> Machida, Ogawa et Okiyama, 1982 (<i>Metaleptus</i>)	N – 715
<i>radiatum</i> Dujardin, 1845 (<i>Paryphostomum</i>)	D – 207
<i>radiatum</i> (Loos, 1900) (<i>Cylicocycclus</i>)	N – 249
<i>radiatum</i> (Rudolphi, 1803) (<i>Oesophagostomum</i>)	N – 537
<i>cf.ralli</i> Underwood et Dronen, 1986 of Scholz and Salgado-Maldonado,(2001) (<i>Cycluster</i>)	C – 312
<i>ralli</i> larv. (Underwood et Dronen, 1986) (<i>Cycluster</i>)	C - 399
<i>ralli</i> Baylis, 1936 (<i>Notocotylus</i>)	D – 185
<i>ramulosus</i> Malewitszkaja, 1941 (<i>Dactylogyrus</i>)	M – 49
<i>ranae</i> (Frölich, 1791) (<i>Opisthioglyphe</i>)	D – 58
<i>ranae</i> Morishita, 1926 (<i>Spinitectus</i>)	N – 165
<i>ranae</i> (Schrank, 1788) (<i>Acanthocephalus</i>)	A – 5
<i>ransomi</i> Schultz, 1940 (<i>Diorchis</i>)	C – 122
<i>raphiodoni</i> Moravec, Kohn et Fernandes, 1993 (<i>Guyanema</i>)	N – 595
<i>rarissimus</i> Gussev, 1966 (<i>Dactylogyrus</i>)	M – 19
<i>rarus</i> Wegener, 1910 (<i>Gyrodactylus</i>)	M – 243
<i>rasini</i> Lucký, 1973 (<i>Gyrodactylus</i>)	M – 345
<i>rastellus</i> (Olsson, 1876) (<i>Dolichosaccus</i>)	D - 403
<i>rebecae</i> (Andrade-Salas, Pineda-López et García-Magaña, 1994) (<i>Procamallanus</i>)	N – 663
<i>rectangulata</i> Fuhrmann, 1908 (<i>Biuterina</i>)	C – 79
<i>rectangulum</i> (Bloch, 1782) (<i>Bathybothrium</i>)	C – 17
<i>recurvatum</i> (Linstow, 1873) (<i>Echinoparyphium</i>)	D –196
<i>recurvum</i> (Solger, 1877) (<i>Paratrichosoma</i>)	N – 699
<i>regoi</i> de Chambrier, Scholz et Vaucher, 1996 (<i>Proteocephalus</i>)	C – 245
<i>reniferum</i> (Loos, 1898) (<i>Astiotrema</i>)	D – 16
<i>resecta</i> (Dujardin, 1845) (<i>Baruscapillaria</i>)	N – 775
<i>reticulatus</i> Micelle et Price, 1964 (<i>Urocleidoides</i>)	M – 324
<i>retusus</i> (Dujardin, 1845) (<i>Cephalogonimus</i>)	D – 72
<i>reversati</i> Pariselle et Euzet, 2003	M - 368
<i>revoluta</i> (Rudolphi, 1819) (<i>Chevreuria</i>)	N – 420
<i>revolutum</i> (Froelich, 1802) (<i>Echinostoma</i>)	D – 130
<i>rhacodes</i> (Deardorff et Overstreet, 1978) (<i>Hysterothylacium</i>)	N – 832
<i>rhamdiae</i> Holcman-Spector et Mañé Garzón, 1988 (<i>Proteocephalus</i>)	C – 342
<i>rhamdiae</i> Moravec, Gonzáles Solis et Vargas Vázquez, 1995 (<i>Paracapillaria</i>)	N – 639
<i>rhamphichthydis</i> Moravec, Kohn et Fernandes, 1996 (<i>Cucullanus</i>)	N – 685
“ <i>rhinolophi</i> “ Groschaft (?) (<i>Travassodendrium</i>)	D – 180
<i>rhobalocephala</i> (Reihm, 1881) (<i>Andrya</i>)	C – 387

<i>rhodei</i> Žitňan, 1964 (<i>Gyrodactylus</i>)	M – 157
<i>rigidus</i> (Van Cleave, 1928) (<i>Neoechinorhynchus</i>)	A – 16
<i>rischta</i> Skrjabin, 1917 (<i>Philometra</i>)	N – 69
<i>robertdollfusi</i> Desportes, 1947 (<i>Synhimantus</i>)	N – 485
<i>robusta</i> Szidat, 1928 (<i>Parastrigea</i>)	D – 611
<i>robusta</i> Moravec, Möller et Petersen, 1992 (<i>Philometra</i>)	N – 582
<i>robustum</i> Yamaguti, 1935 (<i>Echinostoma</i>)	D – 128
<i>robustus</i> Malewitszkaja, 1941 (<i>Dactylogyrus</i>)	M – 54
<i>robustus</i> Pichelin, 1999 (<i>Hypoechinorhynchus</i>)	A – 66
<i>rodhaini</i> Brumpt, 1931 (<i>Schistosoma</i>)	D – 380
<i>rodriguesi</i> Baruš, 1968 (<i>Oxyspirura</i>)	N – 292
<i>rogersi</i> Ergens, 1971 (<i>Tetraonchus</i>)	M – 276
<i>rondoni</i> Travassos, 1919 (<i>Rondonia</i>)	N – 579
<i>rosickyi</i> Ryšavý, 1962 (<i>Anomotaenia</i>)	C – 57
<i>rosickyi</i> Ergens, 1970 (<i>Dactylogyrus</i>)	M – 92
<i>rossittensis</i> (Szidat, 1937) (<i>Khawia</i>)	C – 214
<i>rossittensis</i> (Mühling, 1898) (<i>Urotocus</i>)	D – 514
<i>rostrum</i> Gussev, 1955 (<i>Dactylogyrus</i>)	M – 75
<i>rotundata</i> Linstow, 1899 (<i>Hartertia</i>)	N – 306
<i>rotundatus</i> (Rudolphi, 1819) (<i>Acanthocheilus</i>)	N – 716
<i>roytmani</i> Strelkow, 1963 (<i>Tetraonchus</i>)	M – 275
<i>rubra</i> Harwood, 1935 (<i>Strongyluris</i>)	N – 466
<i>rudolphii</i> Moravec, Scholz et Našincová, 1999 (<i>Baruscapillaria</i>)	N – 591
<i>rudolphii</i> Hartwich, 1964 (<i>Contracaecum</i>)	N – 287
<i>rudolphii</i> Hartwich, 1964 (<i>Contracaecum</i>) larvy	N – 860
<i>rugosum</i> (Batsch, 1786) (<i>Eubothrium</i>)	C – 327
<i>rustica</i> Neslobinský, 1911 (<i>Anomotania</i>)	C – 58
<i>rutilensis</i> Gläser, 1974 (<i>Gyrodactylus</i>)	M – 152
<i>rutili</i> (Glazer, 1967) (<i>Paradiplozoon</i>)	M – 299
<i>rutili</i> (Müller, 1780) (<i>Neoechinorhynchus</i>)	A – 15
<i>rutili</i> Gläser, 1965 (<i>Dactylogyrus</i>)	M – 67
<i>rysavyi</i> Baruš et Coy Otero, 1969 (<i>Aplectana</i>)	N – 408
<i>rysavyi</i> Ergens, 1970 (<i>Dactylogyrus</i>)	M – 97
<i>rysavyi</i> Ergens, 1973 (<i>Gyrodactylus</i>)	M – 134
<i>rysavyi</i> Baruš, 1963 (<i>Oxyspirura</i>)	N – 384
<i>rysavyi</i> Sitko, 2004 (<i>Passeritrema</i>)	D – 516
<i>rysavyi</i> Sonin et Baruš, 1971 (<i>Hamatospiculum</i>)	N – 293
<i>saginata</i> Goeze, 1782 (<i>Taenia</i>)	C – 365
<i>saginatus</i> Van Cleave et Bingham, 1949 (<i>Neoechinorhynchus</i>)	A – 11
<i>sagitata</i> (Leuckart, 1842) (<i>Didcocotyle</i>)	M – 126
<i>sagittatus</i> (Kulakovskaja et Akhmerov, 1962) (<i>Atractolytocestus</i>)	C – 340
<i>sagittavaginalis</i> Lim et Furtado, 1984 (<i>Dactylogyrus</i>)	M – 371
<i>sagittus</i> (Grimm, 1872) (<i>Proteocephalus</i>)	C – 33
<i>sahuensis</i> Ling, 1965 (<i>Dactylogyrus</i>)	M – 340
<i>salamandrae</i> (Frölich, 1789) (<i>Brachycoelium</i>)	D – 66
<i>salaris</i> Malmberg, 1953 (<i>Gyrodactylus</i>)	M – 383
<i>salebrosum</i> (Braun, 1901) (<i>Brachydistomum</i>)	D – 510

<i>salgadi</i> Vidal-Martínez, Aguirre-Macedo et Moravec, 1995 (<i>Philometra</i>)	N – 626
<i>salgadoi</i> Caspeta-Mandujano et Moravec, 2000 (<i>Rhabdochona</i>)	N – 765
<i>salmonicola</i> (Ishii, 1916) (<i>Salvelinema</i>)	N – 697
<i>salmonis</i> (Müller, 1780) (<i>Echinorhynchus</i>)	A – 8
<i>saltatrix</i> Ramachandran, 1973 (<i>Philometra</i>)	N – 809
<i>salvelini</i> (Polyansky, 1952) (<i>Pseudocapillaria</i>)	N – 51
<i>salvelini</i> (Schrank, 1790) (<i>Eubothrium</i>)	C – 126
<i>samirae</i> Popazoglo et Boeger, 2000 (<i>Gyrodactylus</i>)	M – 359
<i>sangeri</i> (Cobbold, 1879) (<i>Bathostomum</i>)	N – 478
<i>sanguinea</i> (Rudolphi, 1819) (<i>Philometroides</i>)	N – 74
<i>sanguineum</i> (Sonsino, 1894) (<i>Anchitrema</i>)	D – 100
<i>sapae</i> (Reichenbach-Klinke, 1961) (<i>Paradiplozoon</i>)	M – 303
<i>sarmientoi</i> (Tantaleán, 1974) (<i>Paraeurysorchis</i>)	M – 415
<i>saurotherae</i> (Baruš et Lorenzo Hernandez, 1970) (<i>Baruscapillaria</i>)	N – 300, N – 374
<i>scardiniensis</i> Gläser, 1974 (<i>Gyrodactylus</i>)	M – 149
<i>scardinii</i> Molnár, 1966 (<i>Skrjabillanus</i>)	N – 60
<i>sceleratus</i> (Travassos, 1923) (<i>Parapharyngodon</i>)	N – 361
<i>scelopori</i> (Gedoelst, 1919) (<i>Atractis</i>)	N – 650
<i>scelopori</i> Gedoelst, 1919 (<i>Cyrtosomum</i>)	N – 347
<i>scelopori</i> (Caballero, 1938) (<i>Skrjabinodon</i>)	N – 648
<i>sciaenae</i> Yamaguti, 1941 (<i>Philometra</i>)	N – 703
<i>sclerovaginalis</i> (Lim et Furtado, 1986) (<i>Dactylogyrus</i>)	M – 293
<i>scolecina</i> (Rudolphi, 1819) Hsü, 1935 (<i>Paradilepis</i>)	C – 168
<i>scoliodontis</i> (Baylis, 1931) Johnston et Mawson, 1945 (<i>Terranova</i>)	N – 847
<i>scolopendra</i> (Diesing, 1850) (<i>Schistotaenia</i>)	C – 181
<i>scorpi</i> (Müller, 1776) (<i>Bothriocephalus</i>)	C – 233
<i>seabassi</i> Wu, Li, Zhu et Xie, 2005 (<i>Pseudorhabdosynochus</i>)	M – 405
<i>sebastisci</i> Yamaguti, 1941 (<i>Philometra</i>)	N – 701
<i>secunda</i> (Nicoll, 1906) (<i>Himasthla</i>)	D – 186
<i>sedelnikowi</i> Gvosdev, 1950 (<i>Gyrodactylus</i>)	M – 156
<i>segonzaci</i> Justine, Cassone et Petter, 2002 (<i>Moravecnema</i>)	N – 796
<i>sekulovici</i> Ergens, 1970 (<i>Dactylogyrus</i>)	M – 90
<i>selangoricus</i> Lim, 1987 (<i>Cornudiscoides</i>)	M – 312
<i>semiarmatus</i> (Molin, 1858) (<i>Skrjabinopsolus</i>)	D – 113
<i>semiteres</i> (Zeder, 1800) (<i>Porrocaecum</i>)	N – 193
<i>senticosum</i> (Baylis, 1927) (<i>Nilonema</i>)	N – 852
<i>serata</i> (Frölich, 1789) (<i>Linguatulla</i>)	L – 1
<i>seravschani</i> Osmanov, 1965 (<i>Gyrodactylus</i>)	M – 190
<i>serendipitus</i> Wheeler et Klassen, 1988 (<i>Thylacicleidus</i>)	M – 397
<i>sergeji</i> Ergens, 1988 (<i>Gyrodactylus</i>)	M – 315
<i>serialis</i> (Bovien, 1926) (<i>Bovienia</i>)	C – 353
<i>seriolae</i> (Ishii, 1931) Yamaguti, (<i>Philometroides</i>)	N – 700
<i>seriolae</i> (Yamaguti, 1934) Price, 1939 (<i>Benedenia</i>)	M – 408
<i>serpentina</i> Harwood, 1932 (<i>Amphibiocapillaria</i>)	N – 118
<i>serranellicabrillae</i> Janiszewska, 1949 (<i>Philometra</i>)	N – 854
<i>serrata</i> (Frölich, 1789) (<i>Linguatula</i>)	L – 1
<i>serratospicula</i> Wehr, 1934 (<i>Diplotriaena</i>)	N – 428
<i>serratus</i> (Looss, 1900) (<i>Triodontophorus</i>)	N – 542
<i>serrispatula</i> (Chin, 1963) (<i>Neophyllodistomum</i>)	D – 469
<i>setigera</i> (Frölich, 1789) (<i>Tschertkovilepis</i>)	C – 189

<i>sexalatus</i> Molin, 1860 (<i>Physocephalus</i>)	N – 303
<i>sexalatus</i> (Molin, 1860) (<i>Physocephalus</i>) larvae	N – 666
<i>shattucki</i> Sandground, 1938 (<i>Shattuckins</i>)	N – 311
<i>shikokuensis</i> Moravec, Koudela, Ogawa et Nagasawa, 1998 (<i>Huffmanella</i>)	N – 692
<i>shulmani</i> Ling, 1962 (<i>Gyrodactylus</i>)	M – 198
<i>schejkini</i> Guschanskaja, 1950 (<i>Victorocara</i>)	N – 291
<i>schouteni</i> Moravec et Cambell, 1991 (<i>Huffmanella</i>)	N – 462
<i>schulzi</i> Freitas, Leut et Almeida, 1937 (<i>Aspiculuris</i>)	N – 307
<i>sibirica</i> (Petrov et Gorbunov, 1931) (<i>Physaloptera</i>)	N – 519
<i>sibiricus</i> Ergens, 1980 (<i>Gyrodactylus</i>)	M – 257
<i>sieboldi</i> Leuckart, 1878 (<i>Archigetes</i>)	C – 45
<i>sieboldi</i> Nordmann, 1832 (<i>Ergasilus</i>)	Cr – 8
<i>siluranae</i> Jackson et Tinsley, 1995 (<i>Batrachocamallanus</i>)	N – 828
<i>siluri</i> Osmanov, 1964 (<i>Procamallanus</i>)	N – 15
<i>siluri</i> (Batsch, 1786) (<i>Silurotaenia</i>)	C – 52
<i>simile</i> Moravec et Little, 1988 (<i>Granulinema</i>)	N – 278
<i>similis</i> Wegener, 1910 (<i>Dactylogyrus</i>)	M – 43
<i>similis</i> Moravec et Van As, 2004 (<i>Falcaustra</i>)	N – 819
<i>similis</i> Looss, 1899 (<i>Haematolechus</i>)	D – 402
<i>similis</i> Moravec, Ali et Abdul-Eis, 1991 (<i>Rhabdochona</i>)	N – 564
<i>similis</i> (Mueller, 1936) (<i>Urocleidus</i>)	M – 263
<i>simillimum</i> (Mühling, 1898) (<i>Psilotrema</i>)	D – 284
<i>simoni</i> Rausch, 1949 (<i>Paradilepis</i>)	C – 232
<i>simplex</i> (Rudolphi, 1809) (<i>Anisakis</i>)	N – 78, N – 171,
<i>simplex</i> (cf.) (Rudolphi, 1809) (<i>Anisakis</i>) larv.	N – 79
<i>simplex</i> Kloss, 1966 (<i>Magnivitellinum</i>)	D – 314
<i>simplicimaleata</i> Bychowsky, 1931 (<i>Dactylogyrus</i>)	M – 55
<i>sinensis</i> Ko, 1975 (<i>Echinocephalus</i>)	N – 849
<i>sinensis</i> Hsü, 1935 (<i>Khawia</i>)	C – 46
<i>sinensis</i> Hsü, 1933 (<i>Pingus</i>)	N – 114
<i>singhi</i> Ali, 1957 (<i>Neocamallanus</i>)	N – 150
<i>singularis</i> La Rue, 1911 (<i>Proteocephalus</i>)	C – 34
<i>sinitsini</i> Mueller, 1934 (<i>Plagioporus</i>)	D – 25
<i>siredonis</i> Baird, 1858 (<i>Hedruris</i>)	N – 742
<i>sisowi</i> Skrjabin, 1913 (<i>Tracheophilus</i>)	D – 188
<i>skrjabini</i> Ryšavý, 1955 (<i>Lutztrema</i>)	D – 166
<i>skrjabini</i> (Iwanitzky, 1928) (<i>Crowcrocoecum</i>)	D – 47
<i>skrjabini</i> Mathevossian, 1945 (<i>Hymenolepis</i>)	C – 139
<i>skrjabini</i> Oshmanov, 1940 (<i>Skrjabinopsolus</i>)	D – 125
<i>skrjabini</i> (Spassky et Roytman, 1958) (<i>Tetraonchus</i>)	M – 269
<i>skrjabini</i> (Baskakov, 1924) (<i>Trichocephalus</i>)	N – 338
<i>slovacicus</i> Ergens, 1963 (<i>Gyrodactylus</i>)	M – 186
<i>sobolevi</i> Schaldybin, 1953 (<i>Neoglyphe</i>)	D – 236
<i>sogandaresi</i> Lumsden, 1963 (<i>Saccocoelioides</i>)	D – 316
<i>solidus</i> larv. (Müller, 1776) (<i>Schistocephalus</i>)	C – 375
<i>solidus</i> (Müller, 1776) (<i>Schistocephalus</i>)	C – 429
<i>sommervillae</i> Bray et Gibson, 1995 (<i>Lepidapedon</i>)	D – 545
<i>somnaensis</i> Ergens et Yukhimenko, 1990 (<i>Gyrodactylus</i>)	M – 338
<i>sonsinoi</i> (Linstow, 1894) (<i>Spinicauda</i>)	N – 178
<i>soricina</i> (Baylis, 1934) (<i>Physaloptera</i>)	N – 437

<i>soricis</i> Asakawa, Kamiya et Ohbayashi, 1988 (<i>Glaphyrostomum</i>)	D – 238
<i>soricis</i> Asakawa, Kamiya et Obayashi, 1988 (<i>Aonchotheca</i>)	N – 276
<i>sotoi</i> Caspeta-Mandujano, Cabañas-Carranza et Salgado-Maldonado, 2004 (<i>Spinitectus</i>)	N – 833
<i>southwelli</i> Fuhrmann, 1933 (<i>Nematoparataenia</i>)	C – 162
<i>sphaerula</i> (Rudolphi, 1803) (<i>Strigea</i>)	D – 439
<i>spasskii</i> Tenora et Baruš, 1958 (<i>Armadolepis</i>)	C – 307
<i>spasskyi</i> Strelkow, 1963 (<i>Tetraonchus</i>)	M – 268
<i>spathaceum</i> (Rudolphi, 1819) (<i>Diplostomum</i>)	D – 191
<i>spathans</i> Dietz, 1909 (<i>Drepanocephalus</i>) mtc.	D – 483
<i>spathiger</i> (Railliet, 1896) (<i>Nematodirus</i>)	N – 535
<i>spathulatus</i> Mueller, 1936 (<i>Gyrodactylus</i>)	M – 252
<i>speciosum</i> Monticelli, 1889 (<i>Pelichnibothrium</i>)	C – 268
<i>sphaericus</i> (Klein, 1905) (<i>Pleurogenoides</i>)	D – 230
<i>sphaerocephalus</i> (Rudolphi, 1809) (<i>Cucullanus</i>)	N – 22
<i>sphaerodactyli</i> Baruš et Coy Otero, 1974 (<i>Pharyngodon</i>)	N – 349
<i>sphaerula</i> (Rudolphi, 1803) (<i>Strigea</i>)	D – 89
<i>sphoeroidis</i> Manter, 1947 (<i>Sclerodistomum</i>)	D – 355
<i>sphyrna</i> Linstow, 1878 (<i>Dactylogyrus</i>)	M – 56
<i>spiculigerum</i> (Mühling, 1898) (<i>Psilotrema</i>)	D – 204
<i>spiculogubernaculus</i> Agarwal, 1958 (<i>Procamallanus</i>)	N – 14
<i>spinicauda</i> (Dujardin, 1845) (<i>Pharyngodon</i>)	N – 570
<i>spiniceps</i> (Looss, 1896) (<i>Acanthostomum</i>)	D – 24
<i>spiniferum</i> (La Valette, 1855) (<i>Echinostoma</i>)	D – 289
<i>spinigerum</i> Owen, 1836 (<i>Gnathostoma</i>)	N – 489
<i>spinosa</i> Gendre, 1923 (<i>Cyrnea</i>)	N – 488
<i>spinosissima</i> (Linstow, 1894) (<i>Raillietina</i>)	C – 179
<i>spinosus</i> (Maplestone, 1931) (<i>Spinostromylus</i>)	N – 216
<i>spinulifera</i> (Woodlank, 1935) (<i>Spasskyellina</i>)	C – 258
<i>spinulosa</i> (Cholodkowsky, 1906) (<i>Vigisolepis</i>)	C – 119
<i>spinulosus</i> (Rudolphi, 1809) (<i>Echinochasmus</i>)	D – 570
<i>spirale</i> (Rudolphi, 1795) (<i>Porrocaecum</i>)	N – 401
<i>spiralibursata</i> (Czaplinsky, 1956) (<i>Microsomacanthus</i>)	C – 155
<i>spiralis</i> (Molin, 1858) (<i>Dispharynx</i>)	N – 471
<i>spiralcirra</i> Kohn et Paperna, 1964 (<i>Gussevia</i>)	M - 389
<i>spiropenis</i> Paperna, 1969 (<i>Annulotrema</i>)	M – 327
<i>spiruli</i> Lim et Furtado, 1984 (<i>Dactylogyrus</i>)	M – 368
<i>spratti</i> Moravec et Sey, 1986 (<i>Paracapillaria</i>)	N – 170
<i>sprostonae</i> Ling, 1962 (<i>Gyrodactylus</i>)	M – 218
<i>squaliobarbi</i> Moravec et Sey, 198 (<i>Rhabdochona</i>)	N – 205
<i>aquamata</i> (Linstow, 1883) (<i>Syncuaria</i>)	N – 416
<i>squamata</i> (Linstow, 1883) (<i>Syncuaria</i>) larv.	N – 571
<i>squamatae</i> Harwood, 1932 (<i>Physaloptera</i>)	N – 524
<i>squameus</i> Gussev, 1955 (<i>Dactylogyrus</i>)	M - 393
<i>squamula</i> (Rudolphi, 1819) (<i>Euryhormis</i>)	D – 85
<i>staffordi</i> Tubanguti, 1928 (<i>Glypthelmins</i>)	D – 226
<i>stankovici</i> Ergens, 1970 (<i>Gyrodactylus</i>)	M – 128
<i>stankovici</i> Ergens, 1970 (<i>Dactylogyrus</i>)	M – 89

<i>stefanskii</i> Czaplinski, 1956 (<i>Diorchis</i>)	C – 96
<i>stellorae</i> (Deblock, Biquet et Capron, 1960) (<i>Wardium</i>)	C – 211
<i>stenocephala</i> Railliet, 1884 (<i>Uncinaria</i>)	N – 308
<i>stentorea</i> (Froehlich, 1802) (<i>Anomotaenia</i>)	C – 59
<i>sternae</i> Kulachkova, 1954 (<i>Tetracladium</i>)	D – 152
<i>steudeneri</i> Janicki, 1904 (<i>Rodentolepis</i>)	C – 117
<i>stichospinus</i> (Seamster et Monaco, 1956) (<i>Rhamnocercoides</i>)	M – 419
<i>stoddardi</i> Cram, 1931 (<i>Aproctella</i>)	N – 224
<i>straminea</i> (Goeze, 1782) (<i>Rodentolepis</i>)	C – 118
<i>straminea</i> (Goeze, 1782) (<i>Vampirolepis</i>)	C – 393
<i>strelkovi</i> Ergens et Danilov, 1983 (<i>Gyrodactylus</i>)	M – 130
<i>striatohamus</i> Zhukov, 1981 (<i>Haliotrema</i>)	M – 390
<i>striatum</i> (Zeder, 1800) (<i>Crenosoma</i>)	N – 237
<i>strigis</i> (Yamaguti, 1939) (<i>Brachylecithum</i>)	D – 526
<i>strigis</i> (Frölich, 1802) (<i>Capillaria</i>)	N – 500
<i>strigis</i> (Schrank, 1788) (<i>Strigea</i>)	D – 595
<i>strongylina</i> (Rudolphi, 1819) (<i>Ascarops</i>)	N – 473
<i>strongyluræ</i> (Hopkins, 1954) (<i>Bucephalopsis</i>)	D – 359
<i>strongyluræ</i> (Hopkins, 1954) (<i>Bucephalopsis</i>)	D – 361
<i>strongyluræ</i> Moravec et Ali, 2005 (<i>Philometra</i>)	N – 834
<i>stunkardi</i> (Holl, 1929) (<i>Pisciamphistoma</i>)	D – 245
<i>sturionis</i> Duvernoy, 1842 (<i>Bothrimonus</i>)	C – 382
<i>stylosa</i> (Rudolphi, 1809) (<i>Passerilepis</i>)	C – 173
<i>stylosa</i> (Rudolphia, 1809) (<i>Dicranotaenia</i>)	C – 90
<i>subclavatus</i> (Pallas, 1760) (<i>Diplodiscus</i>)	D – 4
<i>subdolum</i> Jägerskiöld, 1909 (<i>Maritrema</i>)	D – 605
<i>subtilis</i> Pojmanska, 1969 (<i>Leucochloridium</i>)	D – 574
<i>subtriguetrus</i> (Rudolphi, 1814) (<i>Stichorchis</i>)	D – 222
<i>suctorina</i> (Molin, 1860) (<i>Subulura</i>)	N – 431
<i>sudanensis</i> (Saoud, Abdel-Hamid et Ibrahim, 1974) (<i>Allocreadium</i>)	D – 35
<i>sudanensis</i> (McClelland, 1957) (<i>Sandonia</i>)	D – 6
<i>sudobim</i> (Woodland, 1935) (<i>Houssayela</i>)	C – 383
<i>suecicus</i> Nybelin, 1937 (<i>Dactylogyrus</i>)	M – 38
<i>sulcata</i> (Linstow, 1879) (<i>Hymenolepis</i>)	C – 308
<i>sundanensis</i> Lim, 1987 (<i>Cornudiscoides</i>)	M – 310
<i>superbus</i> (Szidat, 1973) Popazoglo et Boeger, 2000 (<i>Gyrodactylus</i>)	M – 356
<i>surmenicola</i> Okada in Dollfus, 1929 (<i>Nybelina</i>)	C – 427
<i>symmetrus</i> (Johnston, 1912) Yamaguti, 1958 (<i>Dolichosaccus</i>)	D – 303
<i>synodontisi</i> Vassiliades, 1973 (<i>Synodontisi</i>)	N – 174
<i>syriaca</i> (Dubois, 1934) (<i>Duboisia</i>)	D – 565
<i>szanagai</i> Ergens, 1971 (<i>Gyrodactylus</i>)	M – 215
<i>tabargan</i> Schulz, 1931 (<i>Ascaris</i>)	N – 365
<i>taeniaeformis</i> (Batsch, 1786) (<i>Taenia</i>)	C – 239
<i>taichui</i> (Nishigori, 1924) (<i>Haplorchis</i>)	D – 260
<i>taichui</i> (Nishigori, 1924) (<i>Haplorchis</i>) mtc.	D – 269
<i>taimeni</i> Ergens, 1971 (<i>Gyrodactylus</i>)	M – 216
<i>takahashii</i> Suzuki in Takahashi, 1929 (<i>Metagonimus</i>) mtc.	D – 553
<i>tasekensis</i> Lim et Furtado, 1986 (<i>Silurodiscoides</i>)	M – 319
<i>tasmanica</i> Johnston et Mawson, 1945 (<i>Capillostrongyloides</i>)	N – 149

<i>tiberianus</i> Paperna, 1960 (<i>Cichlodogyrus</i>)	M - 125
<i>teixeirafreitasi</i> (Caballero, 1971) (<i>Paracapillaria</i>)	N - 631
<i>temperatus</i> (Stafford, 1905) (<i>Megalodiscus</i>)	D - 244
<i>tener</i> (Looss, 1898) (<i>Pleurogenoides</i>)	D - 93
<i>tenerrima</i> (Linstow, 1878) (<i>Paraquimperia</i>)	N - 57
<i>tenuicollis</i> Price, 1935 (<i>Ascocotyle</i>)	D - 344
<i>tenuicollis</i> Price, 1935 (<i>Ascocotyle</i>) mtc.	D - 345
<i>tenuissima</i> (Rudolphi, 1809) (<i>Capillaria</i>)	N - 323
<i>terdentatum</i> (Linstow, 1890) (<i>Megalobatrachonema</i>)	N - 88
<i>teresoides</i> Fuhrmann, 1906 (<i>Hymenolepis</i>)	C - 138
<i>teresoides</i> (Fuhrmann, 1906) (<i>Hamatolepis</i>)	C - 100
<i>tergestina</i> (Stossich, 1883) (<i>Bucephalopsis</i>)	D - 262
<i>tetraphylliformis</i> de Chambrier, Rego et Vaucher, 1999 (<i>Euzetiella</i>)	C - 403
<i>tetrastomus</i> (Rudolphi, 1810) (<i>Proteocephalus</i>)	C - 297
<i>teuthis</i> (MacCallum, 1915) (<i>Haliotrema</i>)	M - 391
<i>texensis</i> Moravec et Huffman, 2000 (<i>Amphibiocapillaria</i>)	N - 752
<i>thaiensis</i> Moravec, Fiala et Dyková, 2004 (<i>Philometra</i>)	N - 827
<i>thecatium</i> (Linton, 1891) Kostylew, 1924 (<i>Leptorhynchoides</i>)	A - 29
<i>thomasi</i> Seibert, 1944 (<i>Diplotriaena</i>)	N - 492
<i>thurstonae</i> Ergens, 1981 (<i>Cichlidogyrus</i>)	M - 124
<i>thymalli</i> (Annenkova-Khlopina, 1923) (<i>Proteocephalus</i>)	C - 50
<i>thymalli</i> Žitňan, 1960 (<i>Gyrodactylus</i>)	M - 153
<i>tiara</i> (Linstow, 1789) (<i>Tanqua</i>)	N - 514
<i>tiberianus</i> Paperna, 1960 (<i>Cichlodogyrus</i>)	M - 125
<i>tibetanus</i> Dzhaliilov, 1980 (<i>Gyrodactylus</i>)	M - 132
<i>tiflophila</i> (Vigueras, 1943) (<i>Helminthoxys</i>)	N - 352
<i>tigrae</i> Rahemo, 1978 (<i>Rhabdochona</i>)	N - 634
<i>tigrinus</i> Mehra et Negi, 1928 (<i>Ganeo</i>)	D - 227
<i>tilapiae</i> (Baylis, 1958) (<i>Acanthosentis</i>)	A - 20
<i>tilapiae</i> Paperna, 1960 (<i>Cichlidogyrus</i>)	M - 122
<i>tincae</i> Shigin et Shigina, 1959 (<i>Skrjabillanus</i>)	N - 59
<i>tincae</i> Malmberg, 1956 (<i>Gyrodactylus</i>)	M - 8
<i>tincae</i> Gussev, 1965 (<i>Dactylogyrus</i>)	M - 17
<i>tincae</i> (Modeer, 1790) (<i>Asymphyllodora</i>)	D - 49
<i>tissensis</i> Zachvatkin, 1951 (<i>Dactylogyrus</i>)	M - 331
<i>tobagoensis</i> Moravec et Kaiser, 1994 (<i>Rhabdias</i>)	N - 619
<i>tonii</i> Ergens, 1970 (<i>Gyrodactylus</i>)	M - 211
<i>tomentosa</i> (Linstow, 1873) (<i>Pseudocapillaria</i>)	N - 55
<i>torquegenitalis</i> Lim et Furtado, 1984 (<i>Dactylogyrus</i>)	M - 370
<i>torta</i> Linton, 1910 (<i>Helicometra</i>)	D - 368
<i>torulosus</i> (Batsch, 1786) (<i>Proteocephalus</i>)	C - 32
<i>trachea</i> (Montagu, 1811) (<i>Syngamus</i>)	N - 397
<i>trachuri</i> Brian, 1903 (<i>Lernanthropus</i>)	Cr - 2
<i>trachyrinci</i> Wayland, Gibson et Sammerville, 1997 (<i>Echinorhynchus</i>)	A - 27
<i>transversale</i> (Rudolphi, 1802) (<i>Allocreadium</i>)	D - 119
<i>transversum</i> (Travassos, 1917) (<i>Lutztrema</i>)	D - 158
<i>transversus</i> (Rudolphi, 1819) (<i>Prosthorhynchus</i>)	A - 42
<i>travassosi</i> (Price, 1938) (<i>Aphanoblastella</i>)	M - 353
<i>travassosi</i> Gutierrez, 1945 (<i>Contraecaecum</i>)	N - 513
<i>travassosi</i> Moravec, Kohn et Fernandes, 1992 (<i>Neoparaseuratum</i>)	N - 578

<i>travassosi paranaensis</i> Moravec, Kohn et Fernandes 1994 (<i>Travassosnema</i>)	N – 621
<i>travnema</i> Pereira, 1938 (<i>Travnema</i>)	N – 584
<i>triacis</i> (Yamaguti, 1952) (<i>Paraorygmatobothrium</i>)	C – 265
<i>triangula</i> (Krabbe, 1869) (<i>Biuterina</i>)	C – 80
<i>tribolodontis</i> Shimazu, 1992 (<i>Allocreadium</i>)	D – 418
<i>tricipola</i> Gibson, 1973 (<i>Pseudanisakis</i>)	N – 30
<i>tricuspis</i> (Fedtsch, 1874) (<i>Diplotriaena</i>)	N – 398
<i>tridentatus</i> (Drasche, 1884) (<i>Camallanus</i>)	N – 528
<i>tridens</i> (Dujardin, 1845) (<i>Capillaria</i>)	N – 223
<i>trifolium</i> (Braun, 1901) (<i>Cladocystis</i>) mtc.	D – 320
<i>trifolium</i> (Linstow, 1905) (<i>Microsomacanthus</i>)	C – 91
<i>trigonocephalum</i> (Rudolphi, 1808) (<i>Bunostomum</i>)	N – 533
<i>trichomycteri</i> Moravec et Shinn, 2004 (<i>Dentinema</i>)	N - 814
<i>triowonis</i> Achmerow, 1952 (<i>Dactylogyrus</i>)	M – 84
<i>triloba</i> (Rudolphi, 1819) (<i>Hysteromorpha</i>)	D – 208
<i>triloba</i> (Rudolphi, 1819) (<i>Hysteromorpha</i>) mtc.	D – 356
<i>triradicatus</i> Lim et Furtado, 1985 (<i>Sundanonchus</i>)	M – 285
<i>triramosum</i> (Yorke et Macfie, 1920) (<i>Cylicocyclus</i>)	N – 248
<i>trispinosa</i> (Leidy, 1852) (<i>Serpinema</i>)	N – 127
<i>trispinosum</i> (Leidy, 1852) (<i>Serpinema</i>)	N – 677
<i>trispinosum</i> (Leidy, 1852) (<i>Serpinema</i>) larv.	N – 683
<i>tritoniispunctati</i> (Diesing, 1851) (<i>Amphibiocapillaria</i>)	N – 91
<i>trituri</i> Grabda, 1959 (<i>Astiotrema</i>)	D – 71
<i>tropica</i> (Manter, 1936) (<i>Genarchella</i>)	D – 310
<i>tropica</i> Mendoza-Franco, Scholz et Cabañas-Caranza, 2002 (<i>Guavinella</i>)	M – 377
<i>truncata</i> Creplin, 1825 (<i>Hadjelia</i>)	N – 310
<i>truncatus</i> (Rudolphi, 1814) (<i>Camallanus</i>)	N – 81
<i>truncatus</i> (Pallas, 1781) (<i>Cyathocephalus</i>)	C – 13
<i>truncatus</i> (Pallas, 1776) (<i>Cyathocephalus</i>) larv.	C – 246
<i>truttae</i> Sinzar, 1955 (<i>Dentitruncus</i>)	A – 37
<i>truttae</i> Schrank, 1788 (<i>Echinorhynchus</i>)	A – 6
<i>truttae</i> Schrank, 1788 larv. (<i>Echinorhynchus</i>)	A - 76
<i>truttae</i> Fabricius, 1794 (<i>Cucullanus</i>)	N – 66
<i>truttae</i> Gläser, 1974 (<i>Gyrodactylus</i>)	M – 166
<i>tuba</i> Linstow, 1878 (<i>Dactylogyrus</i>)	M – 37
<i>tuberculata</i> (Rudolphi, 1819) (<i>Oochoristica</i>)	C – 22
<i>tubicirrus magnus</i> Paperna et Thurston, 1969 (<i>Cichlidogyrus</i>)	M – 123
<i>tulensis</i> Ergens, 1988 (<i>Gyrodactylus</i>)	M – 316
<i>tumidocollis</i> Wagner, 1953 (<i>Proteocephalus</i>)	C – 250
<i>tumidus</i> Van Cleave et Bangham, 1949 (<i>Neoechinorhynchus</i>)	A – 14
<i>turdia</i> (Ku, 1938) (<i>Lyperosomum</i>)	D – 167
<i>turgida</i> Rudolphi, 1819 (<i>Physaloptera</i>)	N – 259
<i>tuvensis</i> Krasnolobova et Timofeeva, 1968 (<i>Platynosomum</i>)	D – 518
<i>tylosuri</i> Moravec et Ali, 2005 (<i>Philometra</i>)	N – 835
<i>ujati</i> Pineda-López et Andrade-Salas, 1989 (<i>Cichlasotrema</i>)	D – 322
<i>undula</i> (Schrank, 1788) Davies, 1935 (<i>Dilepis</i>)	C – 207
<i>unicus</i> Szidat, 1943 (<i>Palaeorchis</i>)	D – 124
<i>unelen</i> Ostrowski de Nüñez, Burgni et Flores, 2003 (<i>Prosterotestes</i>)	D – 470
<i>urceus</i> (Wedl, 1855) (<i>Paradilepis</i>)	C – 315
<i>urna</i> (Grube et Wagener, 1852) (<i>Gyrococtyle</i>)	C – 426

<i>vakhanica</i> Erhardová-Kotrlá, 1970 (<i>Cephaluris</i>)	N – 312
<i>vanbenedenii</i> (Parona et Perugia, 1890) (<i>Ancyrocephalus</i>)	M – 102
<i>vandenbrokæ</i> Dubois, 1970 (<i>Strigea</i>)	D – 617
<i>vanelli</i> (Rudolphi, 1819) (<i>Uvitellina</i>)	D – 618
<i>vaneyi</i> (Tseng, 1930) (<i>Dollfustrema</i>)	D – 466
<i>varani</i> (Stossich, 1895) (<i>Panceriella</i>)	C – 380
<i>variabilis</i> Mizelle et Kritsky, 1969 (<i>Urocleidoides</i>)	M – 280
<i>varicorhini</i> Ergens et Ibragimov, 1976 (<i>Gyrodactylus</i>)	M – 224
<i>varicus</i> Achmerow, 1952 (<i>Ancylo-discoides</i>)	M – 112
<i>varicus</i> (Müller, 1774) (<i>Derogenes</i>)	D – 538
<i>variglandis</i> (Miller et Northup 1926) (<i>Austro-bilharzia</i>)	D – 449
<i>variolæ</i> Moravec, Justine et Rigby, 2006 (<i>Procamallanus</i>)	N – 864
<i>vastator</i> Nybelin, 1924 (<i>Dactylogyrus</i>)	M – 41
<i>vazi</i> Travassos, 1933 (<i>Micropleura</i>)	N – 801
<i>venezuelensis</i> Slimane, Guerrero et Durette-Desset, 1996 (<i>Oswaldocruzia</i>)	N – 798
<i>ventricosum</i> (Rudolphi, 1809) (<i>Brachylecithum</i>)	D – 431
<i>venusta</i> Freitas et Mondonça, 1958 (<i>Capillaria</i>)	N – 326
<i>venusta</i> (Rosseter, 1897) Spasskaja, 1966 (<i>Retinometra</i>)	C – 97
<i>verai</i> Lamothe-Argumedo et Pireda-López, 1990 (<i>Tabascotrema</i>)	D – 429
<i>vermicularis</i> (Linnaeus, 1758) (<i>Enterobius</i>)	N – 822
<i>verrucosa</i> (Froelich, 1789) (<i>Catatropis</i>)	D – 139
<i>vespertilionis</i> (Müller, 1784) (<i>Plagiorchis</i>)	D – 218
<i>viannai</i> Travassos, 1949 (<i>Cosmoxyrema</i>)	N – 587
<i>vicinoides</i> Ergens et Karimov, 1988 (<i>Gyrodactylus</i>)	M – 329
<i>vicinus</i> Bychowsky, 1957 (<i>Gyrodactylus</i>)	M – 232
<i>vidali</i> González-Solís et Moravec, 2002 (<i>Atractis</i>)	N – 784
<i>vietnamensis</i> Moravec et Sey, 1989 (<i>Batrachotrema</i>)	D – 228
<i>vietnamensis</i> Moravec et Sey, 1987 (<i>Hykynema</i>)	N – 209
<i>vietnamensis</i> Moravec et Sey, 1985 (<i>Omeia</i>)	N – 161
<i>vietnamensis</i> Moravec et Sey, 1988 (<i>Rhabdochona</i>)	N – 204
<i>vietnamensis</i> Moravec et Sey, 1989 (<i>Pseudorhadinorhynchus</i>)	A – 33
<i>vietnamensis</i> Moravec et Sey, 1989 (<i>Prosorhynchus</i>)	D – 224
<i>viguerasi</i> Baruš et Moravec, 1967 (<i>Perezitrema</i>)	D – 12
<i>villosa</i> (Bloch, 1782) (<i>Hymenolepis</i>)	C – 140
<i>vimbi</i> Schulman, 1953 (<i>Gyrodactylus</i>)	M – 214
<i>violentum</i> Van Cleave, 1928 (<i>Hebesoma</i>)	A – 71
<i>viperis</i> (Beddard, 1913) (<i>Ophiotaenia</i>)	C – 230
<i>vistulæ</i> Prost, 1957 (<i>Dactylogyrus</i>)	M – 45
<i>vistulensis</i> (Sivak, 1932) (<i>Ancylo-discoides</i>)	M – 115
<i>vitellioba</i> (Olsson, 1876) (<i>Gorgoderina</i>)	D – 56
<i>viverrini</i> (Poirier, 1886) (<i>Opisthorchis</i>)	D – 254
<i>viverrini</i> (Poirier, 1886) (<i>Opisthorchis</i>) mtc.	D – 270
<i>vivianæ</i> Groschaft et Valle, 1969 (<i>Lecithodendrium</i>)	D – 162
<i>vivipara</i> Wedl, 1862 (<i>Tachygonetria</i>)	N – 410
<i>viviparus</i> (Bloch, 1782) (<i>Dictyocaulus</i>)	N – 540
<i>viviparus</i> (Olsson, 1868) (<i>Zoogonoides</i>)	D – 484
<i>vogtianum</i> Baudon, 1881 (<i>Leucochloridium</i>)	D – 575
<i>vojteki</i> Moravec, Baruš et Ryšavý, 1987 (<i>Spauligodon</i>)	N – 176
<i>volvens</i> Nordmann, 1832 (<i>Diplostomum</i>)	D – 564

<i>vranoviensis</i> Ergens, 1956 (<i>Dactylogyrus</i>)	M – 60
<i>vulgaris</i> (Looss, 1900) (<i>Delafondia</i>)	N – 546
<i>wagneri</i> Nybelin, 1922 (<i>Monobothrium</i>)	C – 206
<i>wagneri</i> Kulwiec, 1927 (<i>Dactylogyrus</i>)	M – 40
<i>walkeri</i> (Ekbaum, 1935) (<i>Salvelinema</i>)	N – 64
<i>wardi</i> Mackin, 1936 (<i>Spironoura</i>)	N – 270
<i>websteri</i> Voge et Read, 1954 (<i>Parafimbriaria</i>)	C – 169
<i>whorligenitalis</i> (<i>Dactylogyrus</i>)	M – 292
<i>willeyi</i> Linstow, 1904 (<i>Kalicephalus</i>)	N – 242
<i>williamsi</i> Bray et Gibson, 1988 (<i>Paralepidapedon</i>)	D – 546
<i>wilsoni</i> Shipley, 1907 (<i>Diphyllobothrium</i>)	C – 330
<i>winteri</i> Tores et Soto, 2003 (<i>Hysterothylacium</i>)	N - 803
<i>wislockii</i> (Sandground, 1938) (<i>Vampirolepis</i>)	C – 190
<i>wisniewskii</i> (Ślusarski, 1958) (<i>Nicolla</i>)	D – 246
<i>wunderi</i> Bychowsky, 1931 (<i>Dactylogyrus</i>)	M – 28
<i>xanthocephalum</i> Monticelli, 1892 (<i>Ceratobothrium</i>)	C – 316
<i>xanthosomus</i> (Creplin, 1846) (<i>Metorchis</i>)	D – 293
<i>xenopi</i> Moravec et Cosgrove, 1982 (<i>Pseudocapillaroides</i>)	N – 136
<i>yagii</i> Asakawa, Kamiya et Ohbayashi, 1988 (<i>Stephanskostrongylus</i>)	N – 448
<i>yamagutii</i> Dollfus, 1960 (<i>Nybelinia</i>) larv.	C – 289
<i>yamagutii</i> Martin, 1974 (<i>Paracardicoloides</i>)	D – 499
<i>yamagutii</i> Martin, 1974 larvae (<i>Paracardicoloides</i>)	D – 500
<i>yangtzensis</i> Yu et Wu, 1989 (<i>Brentisentis</i>)	A – 73
<i>yokogawai</i> Katsurada, 1912 (<i>Metagonimus</i>)	D – 422
<i>yokogawai</i> (Katsuta, 1932) (<i>Haplorchis</i>)	D – 257
<i>yucatanense</i> Moravec, Vidal-Martínez, Vargas.Vázquez, Vivas-Rodríguez, González-Solís, Mendoza-Franco, Simá-Alvarez et Güemez-Ricalde, 1997 (<i>Allonematobothrium</i>)	D – 376
<i>yucatanensis</i> Moravec, Scholz et Vivas-Rodríguez, 1995 (<i>Pseudocapillaria</i>)	N – 642
<i>yucatanensis</i> Vidal-Martínez, Aguirre-Macedo et Mendoza-Franco, 1997 (<i>Pseudorhabdosynochus</i>)	M – 346
<i>yukhimenkoi</i> Ergens, 1978 (<i>Gyrodactylus</i>)	M – 241
<i>zabensis</i> Amin, Abdullah et Mhaisen, 2003 (<i>Neoechinorhynchus</i>)	A - 77
<i>zacconis</i> Yamaguti, 1935 (<i>Rhabdochona</i>)	N – 38
<i>zakharovi</i> Skrjabin, 1920 (<i>Eucotyle</i>)	D – 197
<i>zandti</i> Bychowsky, 1933 (<i>Dactylogyrus</i>)	M – 29
<i>zarudnyi</i> (Skrjabin, 1924) (<i>Tamerlania</i>)	D - 478
<i>zweifeli</i> Moravec et Sey, 1986 (<i>Aplectana</i>)	N – 168
<i>zweifeli</i> Moravec et Sey, 1989 (<i>Halipegus</i>)	D – 235

