

Preliminary Insights into the Impact of Dietary Starch on the Ciliate, *Neobalantidium coli*, in Captive Chimpanzees

Kateřina Schovancová¹*, Kateřina Pomajbíková^{2,4,*,9}, Petr Procházka³, David Modrý^{2,4,5}, Petra Bolechová^{6,7}, Klára J. Petřelková^{2,3,4,7}

1 Institute of Botany and Zoology, Masaryk University, Brno, Czech Republic, **2** Department of Pathology and Parasitology, University of Veterinary and Pharmaceutical Sciences, Brno, Czech Republic, **3** Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, Brno, Czech Republic, **4** Biology Center of the Academy of Sciences of the Czech Republic, Institute of Parasitology, České Budějovice, Czech Republic, **5** Central European Institute for Technology, Brno, Czech Republic, **6** Department of Husbandry and Ethology of Animals, CULS, Prague, Czech Republic, **7** Liberec Zoo, Liberec, Czech Republic

Abstract

Infections caused by the intestinal ciliate *Neobalantidium coli* are asymptomatic in most hosts. In humans and captive African great apes clinical infections occasionally occur, manifested mainly by dysentery; however, factors responsible for development of clinical balantidiasis have not been fully clarified. We studied the effect of dietary starch on the intensities of infection by *N. coli* in two groups of captive chimpanzees. Adult chimpanzees infected by *N. coli* from the Hodonín Zoo and from the Brno Zoo, Czech Republic, were fed with a high starch diet (HSD) (average 14.7% of starch) for 14 days, followed by a five-day transition period and subsequently with a period of low starch diet (LoSD) (average 0.1% of starch) for another 14 days. We collected fecal samples during the last seven days of HSD and LoSD and fixed them in 10% formalin. We quantified trophozoites of *N. coli* using the FLOTAC method. The numbers of *N. coli* trophozoites were higher during the HSD (mean \pm SD: 49.0 \pm 134.7) than during the LoSD (3.5 \pm 6.8). A generalized linear mixed-effects model revealed significantly lower numbers of the *N. coli* trophozoites in the feces during the LoSD period in comparison to the HSD period (treatment contrast LoSD vs. HSD: 2.7 \pm 0.06 (SE), $z = 47.7$; $p < < 0.001$). We conclude that our data provide a first indication that starch-rich diet might be responsible for high intensities of infection of *N. coli* in captive individuals and might predispose them for clinically manifested balantidiasis. We discuss the potential nutritional modifications to host diets that can be implemented in part to control *N. coli* infections.

Citation: Schovancová K, Pomajbíková K, Procházka P, Modrý D, Bolechová P, et al. (2013) Preliminary Insights into the Impact of Dietary Starch on the Ciliate, *Neobalantidium coli*, in Captive Chimpanzees. PLoS ONE 8(11): e81374. doi:10.1371/journal.pone.0081374

Editor: Niko Speybroeck, Université Catholique de Louvain, Belgium

Received: March 31, 2013; **Accepted:** October 13, 2013; **Published:** November 25, 2013

Copyright: © 2013 Schovancová et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the project “CEITEC” – Central European Institute of Technology” (grant no. CZ.1.05/1.1.00/02.0068) from the European Regional Development Fund, by grant from the Grant Agency of the Czech Republic (grant no. 206/09/0927), and by institutional support of Institute of Vertebrate Biology Academy of Sciences of the Czech Republic (grant no. RVO:68081766) and by OPVK 2.3 project – Development of Scientific Team and Laboratory for Infectious Diseases Common to Humans and Great Apes (CZ.1.07/2.3.00/20.0300). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: K. J. Petřelková and P. Bolechová are employed by a commercial company (Liberec Zoo). However, this does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: kpomajbik@gmail.com

⁹ These authors contributed equally to this work.

Introduction

Balantidiasis is a neglected protozoan infection caused by trichostomatid ciliate *Neobalantidium coli* [1] (formerly *Balantidium coli*), inhabiting the hindgut of a variety of mammalian species worldwide [2]. Domestic swine and wild boars are considered as the principle natural reservoirs [3]. In humans, balantidiasis is considered as a zoonotic disease acquired via the fecal-oral route from the reservoir host, usually through contact with swine feces. *N. coli* infection is commonly reported in captive African great apes (prevalence 71%, [4]), but is rarely found in wild apes when examined [4,5].

The factors affecting the pathogenicity of *N. coli* are unclear. It is considered a harmless commensal in reservoir hosts [2], however, in humans and captive African great apes, especially gorillas, the infections can be manifested by a range of clinical symptoms including chronic diarrhea to bloody dysentery [2,6,7,8,9]. Moreover, *N. coli* is able to invade other organs in

immunocompromised human patients [10,11,12,13] and albeit rare, lethal cases have been reported in humans and captive apes [9,14,15,16].

Although pigs and rats have been used to study *Neobalantidium coli* infections [17,18,19], captive African great apes offer a valuable model, which might help us to unravel the factors contributing to the development of clinical balantidiasis in humans, because both humans and African great apes are not natural hosts for *N. coli* [4] and unlike in pigs and rats clinical outcomes have been observed in apes [2,6,7,8,9]. The occurrence of *N. coli* in captive African great apes has been explained by (i) the presence of reservoir(s) in captive facilities or/and (ii) differences in the diet of wild and captive apes [4]. It seems that the reservoir hypothesis alone cannot sufficiently clarify the situation as the suid reservoir hosts occur also in the wild [e.g. red river hogs (*Potamochoerus porcus*) being in close contact with free ranging apes, Pomajbíková, unpublished data]. On the other hand, it is well-known that the diet of captive great apes significantly differs from that in the wild.

Captive ape diets are low in fiber, rich in starch and readily available sugars [20,21,22,23]. Intestinal ciliates from the subclass Trichostomatia are known to possess amylases enabling them to utilize starch, and *in vitro* studies on *N. coli* demonstrated that this ciliate is able to ingest the starch particles [24,25]. Moreover, a handful of very early experimental studies have suggested a possible effect of a starch-rich diet on *N. coli* infection in pigs, rats and humans [17,18,19,26]. In addition, dietary starch has been shown to enhance ciliate growth in a mutualistic entodiniomorphid ciliate *Troglodytella abraxarti* in captive chimpanzees [27].

Our long-term research on *N. coli* in captive primates [1,4] provided us with an opportunity to examine the effect dietary starch on the fecal populations of *N. coli* in two groups of captive chimpanzees.

Materials and Methods

Ethical statement

This study was authorized by managements of Hodonín and Brno Zoos, the ethics committee of the Institute of Vertebrate Biology; AS CR reviewed and approved the protocol of the study. Our research and the protocol complied with the legal requirements of the Czech Republic (Czech National Council Act No. 246/1992 Coll. the protection of animals against cruelty, amended by Act No. 162/1993 Coll.). Our study was non-invasive. Collection of fecal samples did not include any disturbance of the animals (see the part Sample collection and ciliate quantification). We designed the diets in close collaboration with both Zoos. For welfare and enrichment purposes we paid careful attention to feed chimpanzees variable diets of different food components each day. However, in order to design diets which differ in starch, but stay as much as possible similar in all other nutrients, these diets did not fully meet nutritional requirements of chimpanzees in captivity [21,28], but we kept specific feeding habits of each chimpanzee group.

Animals and housing

The study was conducted in the Hodonín Zoo, Czech Republic (chimpanzee group I: adult male Ob, ca. 31 years of age; adult female Zuzana, ca. 37 years of age) from August 12, 2010 to September 19, 2010 and in Brno Zoo, Czech Republic (chimpanzee group II: adult male Fábén, 31 years of age; adult female Nimba, 31 years of age) from November 8, 2010 to December 16, 2010. Chimpanzees in Brno Zoo could utilize three recently repaired indoor enclosures and two outdoor enclosures, while chimpanzees in Hodonín Zoo could use one indoor and one outdoor enclosure. Wooden wool was provided to the chimpanzees for bedding and the indoor enclosures were cleaned once per day by water and disinfectants. Both zoos carry out enrichment programs with focus on feeding enrichment i.e. different ways of hiding, presenting or enriching food provided to animals. Moreover, enclosures are enriched with species specific equipment promoting their natural way of life, including locomotion, resting etc. Chimpanzee had no close contact with humans or other animals kept in the zoos. Our previous study showed that the chimpanzees harbor *N. coli* with no clinical symptoms (e.g. diarrhea) [4].

Diets and sample collection

During the study, we fed the chimpanzee groups two different diets: a high-starch diet (HSD, average 14.7% of starch) and a low-starch diet (LoSD, average 0.1% of starch). The LoSD was composed only from low starch food items, while the HSD included also potatoes and rice (Table 1). First, we fed

chimpanzees the HSD to adapt them to this diet for 7 days, which was followed by a 7-day HSD period with fecal collection. Afterwards there was a 5-day gradual transition to LoSD and another 7-day adaptation LoSD period followed by a 7-day LoSD experimental period when we collected the feces. Throughout the study, daily food intake was 4.502 ± 0.5 kg/animal in wet weight in both zoos. We weighed each dietary item before feeding and, if any of the items remained, after feeding, in order to quantify the exact amount of the total diet ingested.

We froze the samples of each food item and analyzed them for crude protein, lipid, starch, ash, neutral detergent fiber (NDF), acid detergent fiber (ADF), and acid detergent lignin (ADL), by standard methods approved by the Association of Official Analytical Chemists International [29]. Starch was hydrolyzed with hydrochloric acid and analyzed as free sugars by polarimetry, whereas proteins were determined by using the Carres agents. The resulting values were corrected using optically active compounds dissolved in an ethanol–water mixture [29]. Hemicellulose was calculated as NDF–ADF. Cellulose was calculated as ADF–ADL. LoSD and HSD did not differ in terms protein, fiber and lipid concentrations (Table 2).

Sample collection and ciliate quantification

In both zoos we collected fecal samples 2–3 times per day from individuals in indoor and outdoor enclosures. We did not pool the samples. The collection of fecal samples was non-invasive and did not cause any disturbance to the animals. We could not assign fecal samples to particular individuals as the ethical considerations prohibited us from separating the animals for the duration of the study.

We immediately placed 5 g of feces in 20 ml of 10% formalin and then transferred the samples to the laboratory of the Department of Pathology and Parasitology of the University of Veterinary and Pharmaceutical Sciences Brno for examination. We quantified the trophozoites of *N. coli* under light microscope (Olympus CX 41) with magnification 400× and by FLOTAC 400 apparatus according to the FLOTAC technique protocol [30] optimized for *Neobalantidium* trophozoite quantification using the FS3 Zinc Sulphate flotation solution ($ZnSO_4 \cdot 7H_2O$, s.g. 1.2), [31].

Statistical analyses

We tested the effect of diet on the number of *N. coli* trophozoites present in feces by a generalized mixed-effects model (GLMM) with Poisson error distribution and log link function. Zoo (Hodonin, Brno) was considered as the experimental unit. The numbers of trophozoites (count data) entered the model as the response variable, while the type of diet was a categorical predictor with two levels (HSD and LoSD) and zoo (Hodonín and Brno) a random effect. The GLMM was fitted in the package lme4 [32], using Laplace approximation for parameter estimations [33]. All statistical analyses were conducted in R 2.13.1 [34].

Results

We examined 58 samples during HSD and 67 samples during LoSD from group I, and 39 samples during HSD and 28 samples during LoSD from group II. We found only trophozoites of *N. coli*; no other parasites were present. No clinical symptoms were observed throughout the whole study period. The numbers of *N. coli* trophozoites were higher during the HSD (mean \pm SD: 49.0 ± 134.7) than during the LoSD (3.5 ± 6.8 ; Fig. 1). The generalized linear mixed-effects model showed a significant difference in the numbers of *N. coli* trophozoites detected in

Table 1. Daily intake of experimental diets fed to chimpanzees (means of wet matter, g/day and chimpanzee).

	Hodonín		Brno	
	HSD	LoSD	HSD	LoSD
fruit – both diets ¹	2343±196	2283±245	1085±134	1122±251
vegetable – both diets ²	1613±228	1739±259	1862±203	1907±225
vegetable/fruit – only LoSD ³	0	953±108	0	954±171
rice	382±416	0	498±486	0
potatoes (fresh and boiled), rice	618±416	0	428±475	0
dairy products (cottage cheese, yoghurt)	15±42	15±42	92±114	95±118

HSD – high starch diet; LoSD – low starch diet. ¹Zoo Hodonín: orange, grapes, apple, pineapple, banana, kiwi, pears, watermelon, plum, nectarine; ¹Zoo Brno: orange, grapes, apple, pineapple, banana, kiwi; ²both zoos: pepper, tomato, carrot, kohlrabi, cucumber; ³Zoo Hodonín: onion, leek, cauliflower, beet, spring onion, lettuce, Chinese leaves, nectarine; ³Zoo Brno: onion, leek, cauliflower, beet, lettuce, Chinese leaves.

doi:10.1371/journal.pone.0081374.t001

chimpanzee feces between the two diets. The treatment contrast between LoSD and HSD from the GLMM was highly significant (2.7 ± 0.06 (SE), $z = 47.7$; $p < < 0.001$).

Discussion

Early studies, carried out nearly a century ago [17,18,19], demonstrated that high content of starch in the diet of rats and domestic pigs makes the environment in the caecum and large intestine favorable for multiplying of *N. coli* trophozoites, leading to increased infection intensities. In our model situation, which is much closer to humans, we also recorded lower numbers of *Neobalantidium* trophozoites in feces when chimpanzees were fed with the low starch diet. Although the number of experimental units in our study was low, our results provide an indication that starch-rich diet, which is not typical of wild African great apes [21,22,35] might contribute to increased intensities of infection of *N. coli* in captive apes. We point out that a possibility that starch-rich diet might predispose the animals for clinically manifested balantidiasis needs to be further studied.

The assessment of the exact amount of starch passing through the small intestine to the colon of studied chimpanzees is complicated, because resistant starch is determined complexly by several factors and its analytical measurement is difficult [36].

Table 2. Mean nutritional composition of the low- starch diet (LoSD) and high-starch diet (HSD) consumed by chimpanzees.


% of dry matter	Zoo Hodonín		Zoo Brno	
	HSD	LoSD	HSD	LoSD
NDF	10.9±0.5	13.0±0.9	11.0±0.7	12.7±0.6
ADF	8.8±0.4	10.6±1.0	8.3±0.6	9.7±0.5
ADL	1.5±0.1	1.7±0.2	1.7±0.2	1.7±0.2
Cellulose	7.3±0.4	8.9±0.9	6.6±0.5	8.0±0.5
Hemicellulose	2.1±0.2	2.4±0.2	2.7±0.3	3.0±0.3
Crude protein	11.5±2.1	11.9±1.9	8.5±0.6	8.6±0.6
Starch	18.7±2.0	0.1±0.0	10.7±1.9	0.1±0.1
Lipids	2.2±0.5	1.9±0.2	1.6±0.3	1.6±0.1
Ash	5.8±0.3	7.0±0.5	5.2±0.3	5.8±0.3

NDF – neutral detergent fiber; ADF – acid detergent fiber; ADL – acid detergent lignin.

doi:10.1371/journal.pone.0081374.t002

However, studies have shown that the amount of unabsorbed starch entering the colon is directly related to the quantity ingested and that on average 10% of total dietary starch may escape digestion in the small intestine in humans [37,38,39]. Salivary amylase protein levels are ~6–8 times higher in humans than in chimpanzees which suggest that starch digestion in humans is more efficient than that in chimpanzees [40,41]. Thus it is likely that the proportion of starch entering the colon can be even higher in chimpanzees in comparison to humans.

Clinical balantidiasis in humans and great apes is usually treated by administration of antibiotics, though even repeated treatment does not lead to permanent elimination of *N. coli* in infected apes [7,15]. Our results imply that low starch diet can maintain low numbers of trophozoites in the intestine. However, as stated above, our experimental diets designed in order to test the impact of dietary starch did not meet fully nutritional requirements of chimpanzees in captivity (e.g. in protein and fiber content) [21,28] and even the LoSD should not be fed to chimpanzees in long-term. Well-balanced diets, more closely resembling the diet of

**Figure 1.** Numbers of *Neobalantidium coli* in chimpanzee feces during low-starch (LoSD) and high-starch (HSD) diets in two study groups.

doi:10.1371/journal.pone.0081374.g001

wild great apes should be implemented in accordance with modern zoo standards (see [21,22,23,28]). We also stress that due to low number of experimental units, our results need to be considered as preliminary and more research is necessary to look at long-term effects of different starch levels on the number of trophozoites of *N. coli*. Negative impact of starch-rich (and concurrently low-fiber) diet on various components of intestinal microflora of chimpanzees has been described also by other authors ([42] – increase of *Clostridium perfringens*, [43] – occurrence of trichomonads and mycoplasmas, [27,44] – increase of *Troglodytella abrossarti*).

Development of effective antiparasitic strategies requires substantial knowledge of host-parasite interactions, including direct effect of host nutrition on the parasites. To date, most of the research has focused on utilization of nutritional alternations in the control of helminthiasis parasitizing livestock [45,46,47]. Taking into consideration the results of diet experiments with *Neobalantidium* infection in rats and domestic pigs [17,18,19], together with results of our study on chimpanzees, we call for more studies supporting the indications that low starch diet can be implemented as supportive treatment of balantidiasis in addition to antibiotics. In addition, Green and Scully [26] cured four cases of balantidiasis in human patients by putting the patients on milk diet with reduction or complete elimination of starches, while there was no uniformity in dietary management of balantidiasis in other two

studies, which noted diet only incidentally [48,49]. Maybe it is time to reappraise the old studies, describing the impact of diet on *N. coli* infections and future studies should survey the possibilities of nutritional adjustments as a part of parasite control strategies not only in livestock farming, but also in zoo husbandry and potentially in human medicine.

Acknowledgments

We would like to express our sincere thanks to the Hodonín and Brno Zoos for enabling us to conduct this study. We are particularly grateful to Jaroslav Hyjánek, Jiří Ingr, Petra Bilková, Lukáš Baránek (Zoo Hodonín) and Martin Hovorka, Jiří Vítek, Iva Slaninová, Marcela Němečková, Mariana Hubíková (Zoo Brno). We would like to thank to Dr. Jiří Kamler from the Faculty of Forestry and Wood Technology Mendel University in Brno, Czech Republic for his valuable advice regarding nutritional analyses. The manuscript benefited from valuable comments and language revision by Dr. J. M. Rothman from the Department of Anthropology, Hunter College at the City University of New York (USA).

Author Contributions

Conceived and designed the experiments: KS KP PP PB KJP. Performed the experiments: KS KP PB KJP. Analyzed the data: KS KP PP DM KJP. Contributed reagents/materials/analysis tools: KP PP DM PB KJP. Wrote the paper: KS KP PP DM KJP.

References

- Pomajbíková K, Oborník M, Horák A, Petřelková KJ, Grim JN, et al. (2013) Novel insights into the genetic diversity of *Balantidium* and *Balantidium*-like cyst-forming ciliates. *PLoS Negl Trop Dis* 7: e2140.
- Schuster FL, Ramirez-Avila L (2008) Current world status of *Balantidium coli*. *Clin Microbiol Rev* 21: 626–638.
- Solaymani-Mohammadi S, Petri WA jr. (2006) Zoonotic implications of the swine-transmitted protozoal infections. *Vet Parasitol* 140: 189–203.
- Pomajbíková K, Petřelková KJ, Profousová I, Petrášová J, Modrý D (2010) Discrepancies in the occurrence of *Balantidium coli* between wild and captive African great apes. *J Parasitol* 96: 1139–1144.
- Levecke B, Dorny P, Geurden T, Vercammen F, Vercruyse J (2007) Gastrointestinal protozoa in non-human primates of four zoological gardens in Belgium. *Vet Parasitol* 148: 236–246.
- Isidoro J, Martínez J, Ramis A (2013) Haemorrhagic necrotizing colitis associated with *Campylobacter jejuni* and *Balantidium coli* in Zoo gorilla. *J Comp Pathol* 148: 87.
- Lee RV, Prowten AW, Anthone S, Satchidanand SK, Fisher JE, et al. (1990) Typhlitis due to *Balantidium coli* in captive lowland gorillas. *Rev Infect Dis* 12: 1052–1059.
- Teare JA, Loomis MR (1982) Epizootic of balantidiasis in lowland gorillas. *J Am Vet Med Assoc* 181: 1345–1347.
- Kim JCS, Abec CR, Wolf RH (1978) Balantidiosis in a chimpanzee (*Pan troglodytes*). *Lab Anim* 12: 231–233.
- Anargyrou K, Petrikos GL, Suller MT, Skiada A, Siakantaris MP, et al. (2003) Pulmonary *Balantidium coli* infection in a leukemic patient. *Am J Hematol* 73: 180–183.
- Cermeño JR, Hernández De Cuesta I, Uzcátegui O, Páez J, Rivera M, et al. (2003) *Balantidium coli* in a HIV-infected patient with chronic diarrhea. *AIDS* 17: 941–942.
- Clyti E, Aznar C, Couppie P, el Guedj M, Carme B, et al. (1998) A case of co-infection by *Balantidium coli* and HIV in French Guiana. *Bull Soc Pathol Exot* 91: 309–311.
- Ferry T, Bouhour D, De Monbrison F, Laurent F, Dumouchel-Champagne H, et al. (2004) Severe peritonitis due to *Balantidium coli* acquired in France. *Eur J Clin Microbiol Infect Dis* 23: 393–395.
- Dorfman S, Rangel O, Bravo LG (1984) Balantidiasis: report of a fatal case with appendicular and pulmonary involvement. *Trans R Soc Trop Med Hyg* 78: 833–834.
- Lancker F, Mätz-Rensing K, Kiyang J, Jensen SA, Weiss S, et al. (2008) Fatal ulcerative colitis in a western lowland gorilla (*Gorilla gorilla gorilla*). *J Med Primatol* 37: 297–302.
- Pinheiro MC, de Lima MA (1991) A fatal case of intestinal balantidiasis. *Rev Soc Bras Med Trop* 24: 173–176.
- Pritze F (1928) Beiträge zur Kenntnis des *Balantidium coli*. *Ztschr F Parasit* 1: 345–415.
- Schumaker E (1930) *Balantidium coli*: host specificity and relation to the diet of an experimental host. *Am J Epidemiol* 25: 384.
- Schumaker E (1931). Relation of *Balantidium coli* infection to the diet and intestinal flora of the domestic pig. *Am J Hyg* 13: 576–584.
- Schmidt DA, Dempsey JL, Kerley MS, Porton JJ (2000) Fiber in ape diet. A review. In: *Proceedings of the Apes: Challenges for the 21st Century*. Chicago, Illinois: Brookfield Zoo. pp. 177–179.
- NRC (National Research Council) (2003) *Nutrient Requirements of Nonhuman Primates*, 2nd Revised Edition. Washington: The National Academies Press. 286 p.
- Oftedal OT, Allen ME (1996) The feeding and nutrition of omnivores with emphasis on primates. In: Kleiman DG, Allen ME, Thompson KV, Lumpkin S, editors. *Wild mammals in captivity: principles and techniques*. Chicago: University of Chicago Press. pp. 148–157.
- Schwitzer C, Polowinsky SY, Solman C (2009) Fruits as foods – common misconceptions about frugivory. In: Claus M, Fidgett AL, Hatt JM, Huismann T, Hummel J, et al., editors. *Zoo animal nutrition IV*. Fürth: Filander Verlag. pp. 131–168.
- Zaman V (1978) *Balantidium coli*. In: Kreier JP, editor. *Parasitic protozoa*. New York: Academic Press, pp. 633–653.
- Nelson Clifford E (1933) The feeding reactions of *Balantidium coli* from the chimpanzee and pig. *Am J Epidemiol* 18: 185–201.
- Green JJ, Scully FJ (1923) Diet in treatment of *Balantidium coli* infection. *J Am Med Assoc* 81: 291–293.
- Schovanová K (2011) The influence of the diet changes on the populations of intestinal ciliates in captive chimpanzees, master thesis. Brno: Masaryk University. pp. 57.
- AZA Ape TAG (2010) *Chimpanzee (Pan troglodytes) Care Manual*. Silver Spring, MD: Association of Zoos and Aquariums.
- AOAC (2012) *Official methods of analysis of AOAC International*, 18th Edition. Gaithersburg, USA: AOAC International.
- Cringoli G, Rinaldi L, Maurelli MP, Utzinger J (2010) FLOTAC: new multivalent techniques for qualitative and quantitative coproscopic diagnosis of parasites in animals and humans. *Nat Protoc* 5: 503–515.
- Růžicová M (2012) The utilizing of the Flotac® method in the diagnostics of intestinal ciliates, master thesis. Brno: University of Veterinary and Pharmaceutical Sciences Brno. pp. 51.
- Bates D, Maechler M, Dai B (2008) lme4: Linear mixed-effects models using Eigen and R syntax. R package version 0.999375–28. <http://lme4.r-forge.r-project.org/>.
- Bolker BM, Brooks ME, Clark CJ, Geange SW, John R Poulsen, et al. (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends Ecol Evol* 24: 127–135.
- R Development Core Team (2011) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.
- Hockings KJ, Anderson JR, Matsuzawa T (2010) Flexible feeding on cultivated underground storage organs by rainforest-dwelling chimpanzees at Bossou, West Africa. *J Hum Evol* 58: 227–23.

36. McCleary BV, McNally M, Rossiter P (2002) Measurement of resistant starch by enzymatic digestion in starch and selected plant materials: collaborative study. *J AOAC Int* 85: 1103–11.
37. Chapman RW, Sillery JK, Graham MM, Saunders DR (1985) Absorption of starch by healthy ileostomates – effect of transit-time and of carbohydrate load. *Am J Clin Nutr* 41: 1244–1248.
38. Stephen AM, Haddad AC, Phillips SF (1983) Passage of carbohydrate into the colon. Direct measurement in humans. *Gastroenterol* 85: 589–595.
39. Topping DL, Clifton PM (2001) Short chain fatty acids and human colonic function: roles of resistant starch and non-starch polysaccharides. *Physiol Rev* 81: 1031–1064.
40. McGeachin RL, Akin JR (1982) Amylase levels in the tissues and body fluids of several primate species. *Comp Biochem Physiol A* 72: 267–9.
41. Perry GH, Dominy NJ, Claw KG, Lee AS, Fiegler H, et al. (2007) Diet and the evolution of human amylase gene copy number variation. *Nat Genet* 39: 1256–1260.
42. Fujita S, Kageyama T (2007) Polymerase chain reaction detection of *Clostridium perfringens* in feces from captive and wild chimpanzees, *Pan troglodytes*. *J Med Primatol* 36: 25–32.
43. Kisidayova S, Varadyova Z, Pristas P, Piknova M, Nigutova K, et al. (2009) Effects of high- and low-fiber diets on fecal fermentation and fecal microbial populations of captive chimpanzees. *Am J Primatol* 71: 548–557.
44. Petrzalkova KJ, Schovancova K, Profousova I, Kisidayova S, Varadyova Z (2012) The effect of low- and high-fiber diets on the population of entodiniomorphid ciliates *Troglodytella abrossarti* in captive chimpanzees (*Pan troglodytes*). *Am J Primatol* 74: 669–675.
45. Das G, Abel H, Rautenschlein S, Humburg J, Schwarz A, et al. (2011) Effects of dietary non-starch polysaccharides on establishment and fecundity of *Heterakis gallinarum* in grower layers. *Vet Parasitol* 178: 121–128.
46. Houdijk JGM, Kyriazakis I, Kidane A, Athanasiadou S (2012) Manipulating small ruminant parasite epidemiology through the combination of nutritional strategies. *Vet Parasitol* 186: 38–50.
47. Roepstorff A, Mejer H, Nejsum P, Thamsborg SM (2011) Helminth parasites in pigs: New challenges in pig production and current research highlights. *Vet Parasitol* 180: 72–81.
48. Nisbet WO (1920) A case of *Balantidium coli* infection. *South M J* 13: 403.
49. Steen PH (1921) Balantidiasis. *Ugeskr F Laeger* 83: 662.